

6.7. Irodalmi hivatkozások³⁵

Hivatkozások a 2.1. szakaszban

- [1] H.M. Tenney, "Selectivity of various liquid substrates used in gas chromatography", ANAL. CHEM. **30** (1958) 2–8.
- [2] L. Rohrschneider, "Polarity of the stationary phase in gas chromatography", Z. ANAL. CHEM. **170** (1959) 256–263 (CA 54:3045i)³⁶.
- [3] L. Rohrschneider, "Method of characterization for gas chromatographic separation of liquids", J. CHROMATOGR. **22** (1966) 6–22 (CA 65:1421e).
- [4] L. Rohrschneider, in ADVANCES IN CHROMATOGRAPHY, VOL. 4, J.C. Giddings, R.A. Keller, Eds., Marcel Dekker, New York (1967) 333–363.
- [5] W.O. McReynolds, "Characterization of some liquid phases", J. CHROMATOGR. SCI. **8** (1970) 685–691.
- [6] Z. Szentirmai, G. Tarjan, J. Takacs, "Contribution to the theory of the retention index system. Part V. Characterization of polarity of stationary phases. McReynolds system", J. CHROMATOGR. **73** (1972) 11–18.
- [7] L. Rohrschneider, "Chromatographic characterization of liquid phases and solutes for column selection and identification", J. CHROMATOGR. SCI. **11** (1973) 160–166.
- [8] J.J. Leary, J.B. Justice, S. Tsuge, S.R. Lowry, T.L. Isenhour, "Correlating gas chromatographic liquid phases by means of a nearest neighbour technique: a proposed set of twelve preferred phases", J. CHROMATOGR. SCI. **11** (1973) 201–206.
- [9] A. Hartkopf, "A reconsideration of the Rohrschneider approach for characterizing gas chromatographic liquid phases. Part I. Review and description of basic approach", J. CHROMATOGR. SCI. **12** (1974) 113–118.
- [10] L.R. Snyder, "Classification of solvent properties of common liquids", J. CHROMATOGR. SCI. **16** (1978) 223–234.
- [11] C.F. Poole, S.K. Poole, "Characterization of solvent properties of gas chromatographic liquid phases", CHEM. REV. **89** (1989) 377–395.
- [12] T.O. Kollie, C.F. Poole, "Application of multivariate analysis to the selection of test solutes for characterizing stationary phase selectivity in gas chromatography", J. CHROMATOGR. **550** (1990) 213–237.
- [13] C.F. Poole, T.O. Kollie, S.K. Poole, "Recent advances in solvation models for stationary phase characterization and the prediction of retention in gas chromatography", CHROMATOGRAPHIA **34** (1992) 281–302.
- [14] G. Castello, G. D'Amato, "Fast evaluation of the polarity of gas chromatographic columns using the difference in apparent carbon number of linear alkanes and alcohols with the same retention", J. CHROMATOGR. **623** (1992) 289–295.
- [15] L. Rohrschneider, "Development and use of phase constants for the characterization of the stationary phase selectivity in gas chromatography", LC-GC INT. **6** (1993) 553–561.
- [16] G. Castello, G. D'Amato, S. Vezzani, "Evaluation of the polarity of packed and capillary columns by different classification methods", J. CHROMATOGR. **646** (1993) 361–368.
- [17] L. Rohrschneider, "Explanatory coefficients for stationary phases in gas chromatography from McReynolds phase constants", CHROMATOGRAPHIA **38** (1994) 679–688.
- [18] J.A. Yancey, "Review of liquid phases in gas chromatography. Part I. Intermolecular forces", J. CHROMATOGR. SCI. **32** (1994) 349–357.
- [19] J.A. Yancey, "Review of liquid phases in gas chromatography. Part II. Applications", J. CHROMATOGR. SCI. **32** (1994) 403–413.
- [20] S.K. Poole, C.F. Poole, "Chemometric classification of the solvent properties (selectivity) of commonly used gas chromatographic stationary phases", J. CHROMATOGR. A. **697** (1995) 415–427.

³⁵ A dolgozat elkészítéséhez végzett irodalmazást 1998 végén zártam le.

³⁶ A *Chemical Abstract*-re történő hivatkozások formája: CA kötetszám:cikkelyazonosító.

- [21] S.K. Poole, C.F. Poole, "Application of principal component factor analysis to the cavity model of solvation to identify factors important in characterizing the solvent properties of gas chromatographic stationary phases", *J. CHROMATOGR. A.* **697** (1995) 429–440.
- [22] L. Rohrschneider, "Dependence of GC-retention data on phase constants a and b in the James–Martin relationship for homologous n-alkanes", *CHROMATOGRAPHIA* **42** (1996) 489–493.

Hivatkozások a 2.2. szakaszban

- [1] G. Schomburg, "Sampling systems in capillary chromatography", in *PROC. 4TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Bad Duerkheim, FRG, 1981), R.E. Kaiser, Ed., Institute of Chromatography, Bad Duerkheim (1981) 371–404 (CA 97:229484m).
- [2] W. Bertsch, "Methods in high resolution gas chromatography. Two-dimensional techniques. Part 3", *J. HIGH RESOLUT. CHROMATOGR.* **1** (1978) 289–299.
- [3] "The Sadtler standard gas chromatography retention index library", Sadtler Research Laboratories, Philadelphia, Pennsylvania, USA (1986).
- [4] V. Lopez-Avila, J. Benedicto, E. Baldin, W.F. Beckert, "Analysis of classes of compounds of environmental concern. Part I. Nitroaromatic compounds", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 601–607.
- [5] V. Lopez-Avila, J. Benedicto, E. Baldin, W.F. Beckert, "Analysis of compounds of environmental concern. Part I. Diphenyl ethers", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 160–164.
- [6] G. Castello, A. Timossi, T.C. Gerbino, "Gas chromatographic separation of halogenated compounds on non-polar and polar wide bore capillary columns", *J. CHROMATOGR.* **454** (1988) 129–143.
- [7] R.E. Shirey, "Rapid analysis of environmental samples using solid-phase microextraction (SPME) and narrow bore capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **18** (1995) 495–499.
- [8] J.F. Schneider, S. Bourne, A.S. Boparai, "Parallel capillary gas chromatography in the determination of chlorinated pesticides and PCBs", *J. CHROMATOGR. SCI.* **22** (1984) 203–206.
- [9] L. Toretti, A. Simonella, "Multidimensional capillary GC of bromopropylate residues in honey: comparison of some identification procedures", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 142–145.
- [10] L. Torretti, A. Simonella, A. Dossena, E. Torretti, "Determination of organochlorine pesticide residues by solid phase extraction and dual-column HRGC", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 99–101.
- [11] V. Lopez-Avila, J. Benedicto, E. Baldin, W.F. Beckert, "Analysis of classes of compounds of environmental concern. Part III. Organochlorine pesticides", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 319–328.
- [12] Z. Penton, "GC determination of halogenated pesticides with a temperature-programmable injector, dual columns and dual ECDs", in *PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 738–743.
- [13] M.S. Rahman, S. Bøwadt, B. Larsen, "Dual-column GC analysis of Mediterranean fish for 10 organochlorine pesticides and 62 chlorobiphenyls", in *PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 815–821.
- [14] E. Storr-Hansen, "Comparative analysis of thirty polychlorinated biphenyl congeners on two capillary columns of different polarity with non-linear multi-level calibration", *J. CHROMATOGR.* **558** (1991) 375–391.
- [15] E. Storr-Hansen, "Simultaneous analysis for 32 PCB-congeners on two capillary columns operated in parallel with glass T-split", *INT. J. ENVIRON. ANAL. CHEM.* **43** (4) (1991) 253–266 (CA 115: 84564r).
- [16] M.S. Rahman, S. Bøwadt, B. Larsen, "Dual-column GC analysis of mediterranean fish for ten organochlorine pesticides and sixty two chlorobiphenyls", *J. HIGH RESOLUT. CHROMATOGR.* **16** (1993) 731–735.
- [17] S. Bøwadt, B. Johansson, P. Fruekilde, M. Hansen, D. Zilli, B. Larsen, J. de Boer, "Supercritical fluid extraction of polychlorinated biphenyls from lyophilized fish tissue", *J. CHROMATOGR. A.* **675** (1994) 189–204.

- [18] B.R. Larsen, "HRGC separation of PCB congeners", *J. HIGH RESOLUT. CHROMATOGR.* **18** (1995) 141–151.
- [19] P. Hess, J. de Boer, W.P. Cofino, P.E.G. Leonards, D.E. Wells, "Critical review of the analysis of non- and mono-ortho-chlorobiphenyls", *J. CHROMATOGR. A.* **703** (1995) 417–465.
- [20] G.M. Frame, J.W. Cochran, S.S. Bøwadt, "Complete PCB congener distributions for 17 Aroclor mixtures determined by 3 HRGC systems optimized for comprehensive, quantitative, congener-specific analysis", *J. HIGH RESOLUT. CHROMATOGR.* **19** (1996) 657–668.
- [21] G. Castello, S. Vezzani, T.C. Gerbino, "Gas chromatographic separation and automatic identification of complex mixtures of organic solvents in industrial wastes", *J. CHROMATOGR.* **585** (1991) 273–280.
- [22] C.A. Jemma, P.R. Shore, K.A. Widdicombe, "Analysis of C1-C16 hydrocarbons using dual-column capillary GC: application to the exhaust emissions from passenger car and motorcycle engines", *J. CHROMATOGR. SCI.* **33** (1995) 34–48.
- [23] J.M. Levy, "Dual capillary column analysis of complex oxygenated mixtures", *J. CHROMATOGR. SCI.* **22** (1984) 149–152.
- [24] J.M. Levy, J.A. Yancey, "Dual column gas chromatographic analysis of alcohols and methyl tert-butyl ether in gasolines", *J. HIGH RESOLUT. CHROMATOGR.* **9** (1986) 383–387.
- [25] A. D'Amato, C. Bicchi, M. Galli, "The use of three way press-fit connectors (Y connectors) in the analysis of essential oils", *J. HIGH RESOLUT. CHROMATOGR.* **12** (1989) 349–352.
- [26] J.P. Kababick, "Evaluation of incense purity using simultaneous steam distillation-extraction and HRGC analysis", *J. HIGH RESOLUT. CHROMATOGR.* **19** (1996) 241–242.
- [27] V.W. Watts, T.F. Simonick, "Screening of basic drugs in biological samples using dual column capillary chromatography and nitrogen-phosphorus detectors", *J. ANAL. TOXICOL.* **10** (5) (1986) 198–204 (CA 105:204240z).
- [28] I. Ojanperä, I. Rasanen, E. Vuori, "Automated quantitative screening for acidic and neutral drugs in whole blood by dual-column capillary gas chromatography", *J. ANAL. TOXICOL.* **15** (4) (1991) 204–208 (CA 115:152609m).
- [29] C.-W. Oh, J.-H. Kim, K.-R. Kim, D.M. Brownson, T.J. Mabry, "Simultaneous gas chromatographic analysis of non-protein and protein amino acids as N(O,S)-isobutyloxycarbonyl tert.-butyldimethylsilyl derivatives", *J. CHROMATOGR. A.* **669** (1994) 125–137 (CA 121:53281r).
- [30] C.-H. Oh, T.J. Mabry, K.-R. Kim, J.-H. Kim, "GC-MS analysis of nonprotein amino acids in *gymnocladus dioicus* as N(O,S)-isobutyloxycarbonyl silyl derivatives", *J. CHROMATOGR. SCI.* **33** (1995) 399–404.
- [31] C.-H. Oh, J.-H. Kim, K.-R. Kim, T.J. Mabry, "Rapid gas chromatographic screening of edible seeds, nuts and beans for non-protein and protein amino acids", *J. CHROMATOGR.* **708** (1995) 131–141.
- [32] T. Doran, J.P. Cross, "Single sample analysis of the mixture O₂, N₂, CH₄, CO, CO₂, C₂H₆, C₃H₈, and n-C₄H₁₀", *J. GAS CHROMATOGR.* **4** (7) (1966) 260–262 (CA 65:11325b).
- [33] P.K. Gupta, J.G. Nikelly, "Consecutive gas chromatograms from parallel columns using a single injection and a common detector", *ANAL. CHEM.* **63** (1991) 1264–1270.
- [34] L.R. Hogge, D.J.H. Olson, "A simultaneous dual capillary column GC/MS technique for the identification of positional and geometrical isomers of aliphatic compounds", *J. CHROMATOGR. SCI.* **21** (1983) 524–528.
- [35] B.W. Cook, M. Cleary, "Systematic approach to coupled gas chromatography Fourier transform infrared spectroscopy using two wide bore capillary columns", *ANALYST* **113** (1988) 1171–1174.
- [36] J.C. Marr, J. Visentini, M.A. Quilliam, "Parallel column gas chromatography–mass spectrometry for the analysis of complex environmental samples", *INSTRUM. TRACE ORG. MONIT.*, R.E. Clement, K.W.M. Siu, H.H. Hill, Jr., Eds., Lewis, Chelsea, Mich. (1992) 219–240 (CA 116:227289p).
- [37] J.J. Jimenez, J.L. Bernal, J. Atienza, "CGC/AED and CGC/ECD/NPD comparison for the determination of acaricides in honey after hexane/acetone extraction", *CHROMATOGRAPHIA* **42** (1996) 130–134.
- [38] J.L. Bernal, M.J. del Nozal, J. Atienza, J.J. Jimenez, "Multidetermination of PCBs and pesticides by use of a dual GC column–dual detector system", *CHROMATOGRAPHIA* **33** (1992) 67–76.
- [39] G.S. Durell, T.C. Sauer, "Simultaneous dual-column, dual-detector gas chromatographic determination of chlorinated pesticides and polychlorinated biphenyls in environmental samples", *ANAL. CHEM.* **62** (1990) 1867–1871.

- [40] M.L. Hopper, "Analysis of organochlorine pesticide residues using simultaneous injection of two capillary columns with electron capture and electrolytic conductivity detectors", *J. ASSOC. OFF. ANAL. CHEM.* **74** (6) (1991) 974–981 (CA 116:5404h).
- [41] V.M.A. Häkkinen, "Analysis of chemical warfare agents in water by solid phase extraction and two-channel capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 811–815.
- [42] Y. Higashino, "Analysis of components of paint solvent by multidimensional gas chromatography", *SHIKIZAI KYOKAISHI* **62** (12) (1989) 737–743 (CA 113:61347s).
- [43] J. Enqvist, A. Hesso, "Precision gas chromatography as an analytical tool. Part I. a new automatic system based on two-channel retention index monitoring", *KEM.–KEMI* **9** (3) (1982) 176–180 (CA 97:103653x).
- [44] A. Kiviranta, "Retention index method with dual-column high-resolution GC in environmental and food analysis", *INTL. LAB.* **17** (1) (1987) 58–65.
- [45] V.M.A. Häkkinen, "Analysis of chemical warfare agents in water by solid phase extraction and two-channel capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 811–815.
- [46] I. Rasanen, I. Ojanperä, J. Vartiovaara, E. Vuori, P. Sunila, "The advantage of dual-column approach and retention indices combined with refined reporting in gas chromatographic drug screening", *J. HIGH RESOLUT. CHROMATOGR.* **19** (1996) 313–321.
- [47] T. Welsch, U. Dornberger, D. Lerche, "Selectivity tuning of serially coupled columns in high performance liquid chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **16** (1993) 18–26.
- [48] C.J. Little, R.A. Little, M. de la Pena, "A novel design multi-dimensional chromatographic column", *INTL. LABMATE* **19** (7) (1995) 39–41.
- [49] P.H. Lukulay, V.L. McGuffin, "Solvent modulation in liquid chromatography: extension to serially coupled columns", *J. CHROMATOGR. A* **691** (1995) 171–185.
- [50] L. Karlsson, M. Jaremo, M. Emilsson, L. Mathiasson, J.A. Joensson, "Retention in coupled capillary column supercritical fluid chromatography with lipids as model compounds", *CHROMATOGRAPHIA* **37** (1993) 402–410.
- [51] P. Lembke, H. Engelhardt, "Rapid determination of phytanic acid in human blood serum by selectivity tuning in packed column SFC", *J. HIGH RESOLUT. CHROMATOGR.* **16** (1993) 700–702.
- [52] A. Kot, P. Sandra, F. David, "Selectivity tuning in packed column SFC", *J. HIGH RESOLUT. CHROMATOGR.* **17** (1994) 277–279.
- [53] J.A. Rijks, Thesis, University of Technology, Eindhoven, The Netherlands (1973).
- [54] L. Soják, J.A. Rijks, *J. CHROMATOGR.* **119** (1976) 505.
- [55] R. Annino, J. Franko, H. Keller, "Mixed carrier gases in chromatography—A source of error", *ANAL. CHEM.* **43** (1971) 107–117.
- [56] B.A. Rudenko, M.A. Baydarovtseva, V.A. Kuzovkin, V.F. Kucherov, "Gas chromatography with vapours of low-boiling substances as mobile phases", *J. CHROMATOGR.* **104** (1975) 271–275.
- [57] M. Abdel-Rehim, L. Zhang, M. Hassan, H. Ehrsson, "Mixture of ammonia and helium as a carrier gas for capillary gas chromatography", *J. MICROCOLUMN SEP.* **5** (1993) 537–542.
- [58] V. Pretorius, "Mixed mobile phases for controlling relative retention in gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **1** (1978) 199–200.
- [59] K.W.M. Siu, W.A. Aue, "Binary liquid phase system for gas-liquid chromatography with in situ variable polarity", *J. CHROMATOGR.* **189** (1980) 255–258.
- [60] K. Grob, A. Habich, "Effect of water on retention time", *J. HIGH RESOLUT. CHROMATOGR.* **6** (1983) 34–35.
- [61] V.G. Berezkin, A.A. Korolev, "Changes in retention values in open tubular columns under the effect of water vapour" *ZH. ANAL. KHM.* **50** (1995) 1057–1059.
- [62] R.J. Laub, "Variation of gas chromatographic retentions with carrier pressure and composition", *ANAL. CHEM.* **56** (1984) 2115–2119.
- [63] W. Bertsch, "Methods in high resolution gas chromatography, two-dimensional techniques. Part I.", *J. HIGH RESOLUT. CHROMATOGR.* **1** (1978) 85–90.
- [64] W. Bertsch, "Methods in high resolution gas chromatography. two-dimensional techniques. Part II.", *J. HIGH RESOLUT. CHROMATOGR.* **1** (1978) 187–194.
- [65] W. Bertsch, "Methods in high resolution gas chromatography. two-dimensional techniques. Part III.", *J. HIGH RESOLUT. CHROMATOGR.* **1** (1978) 289–299.
- [66] D.R. Deans, "Use of heart cutting in gas chromatography: a review", *J. CHROMATOGR.* **203** (1981) 19–28.

- [67] R.R. Freeman, "Multidimensional gas chromatography", in HIGH RESOLUTION GAS CHROMATOGRAPHY, Second Edition, R.R. Freeman, Ed., Hewlett-Packard (1981) 81–104.
- [68] C.J. Cowper, "Multidimensions for gas chromatography", LABORATORY PRACTICE **34** (1) (1985) 77–80.
- [69] B.M. Gordon, C.E. Rix, M.F. Borgerding, "Comparison of state-of-the-art column switching techniques in high resolution gas chromatography", J. CHROMATOGR. SCI. **23** (1985) 1–10.
- [70] D.E. Willis, "Column switching in gas chromatography", in ADVANCES IN CHROMATOGRAPHY, J.C. Giddings, E. Grushka, P.R. Brown, Eds., **28** (1989) 65–112.
- [71] K. Himberg, E. Sippola, M.L. Riekkola, "Multidimensional gas chromatography: state of the art", J. MICROCOLUMN SEP. **1** (6) (1989) 271–277 (CA 115:105104h).
- [72] W. Bertsch, "Multidimensional gas chromatography", in MULTIDIMENSIONAL CHROMATOGRAPHY TECHNIQUES AND APPLICATIONS, Chromatographic Science Series 50, H.J. Cortes, Ed., Marcel Dekker, New York (1990).
- [73] J.C. Giddings, "Use of multiple dimensions in analytical separation", in MULTIDIMENSIONAL CHROMATOGRAPHY TECHNIQUES AND APPLICATIONS, Chromatographic Science Series 50, H.J. Cortes, Ed., Marcel Dekker, New York (1990).
- [74] U.K. Goekeller, F. Mueller, "Process multidimensional gas chromatography", in MULTIDIMENSIONAL CHROMATOGRAPHY TECHNIQUES AND APPLICATIONS, Chromatographic Science Series 50, H.J. Cortes, Ed., Marcel Dekker, New York (1990).
- [75] L.D. Rothman, "Hardware consideration for the automation of multidimensional chromatographs", in MULTIDIMENSIONAL CHROMATOGRAPHY TECHNIQUES AND APPLICATIONS, Chromatographic Science Series 50, H.J. Cortes, Ed., Marcel Dekker, New York (1990).
- [76] H.J. Cortes, "Developments in multidimensional separation systems", J. CHROMATOGR. **626** (1992) 3–23.
- [77] G. Schomburg, "Two-dimensional gas chromatography: principles, instrumentation, methods", J. CHROMATOGR. A. **703** (1995) 309–325.
- [78] J.B. Phillips, J. Xu, "Comprehensive multi-dimensional gas chromatography ", J. CHROMATOGR. A. **703** (1995) 327–334.
- [79] H.J. Cortes, "Multidimensional chromatography using on-line coupled high-performance liquid chromatography and capillary gas chromatography", in MULTIDIMENSIONAL CHROMATOGRAPHY TECHNIQUES AND APPLICATIONS, Chromatographic Science Series 50, H.J. Cortes, Ed., Marcel Dekker, New York (1990).
- [80] K. Grob, "On-line coupled LC-GC", Hüthig Verlag, Heidelberg (1991).
- [81] J.M. Levy, R.A. Cavalier, T.N. Bosch, A.F. Rynaski, W.E. Huhak, "Multidimensional Supercritical fluid chromatography and supercritical fluid extraction", J. CHROMATOGR. SCI. **27** (1989) 341–346.
- [82] I.L. Davies, K.E. Markides, M.L. Lee, K.D. Bartle, "Multidimensional supercritical fluid chromatography", in MULTIDIMENSIONAL CHROMATOGRAPHY TECHNIQUES AND APPLICATIONS, Chromatographic Science Series 50, H.J. Cortes, Ed., Marcel Dekker, New York (1990).
- [83] L.S. Ette, "Selective detection in column chromatography", J. CHROMATOGR. SCI. **16** (1978) 396–417.
- [84] S.O. Farwell, D.R. Gage, R.A. Kagel, "Current status of prominent selective gas chromatographic detectors: a critical assessment", J. CHROMATOGR. SCI. **19** (1981) 358–376.
- [85] M Dressler, "Selective gas chromatographic detectors", Elsevier, Amsterdam (1986) (CA 106: 148688h).
- [86] R.P.W. Scott, "Chromatographic detectors", Marcel Dekker, New York (1995).
- [87] F.W. Karasek, R.E. Clement, "Basic gas chromatography–mass spectrometry. Principles and techniques", Elsevier, Amsterdam (1988).
- [88] F.A. Settle, M.A. Pleva, "Gas chromatography–mass spectrometry. A knowledge base", Elsevier, Amsterdam (1988).
- [89] N. Ragunathan, K.A. Krock, C. Klawun, T.A. Sasaki, C.L. Wilkins, "Multispectral detection for gas chromatography", J. CHROMATOGR. A. **703** (1995) 335–382.
- [90] P.R. Griffiths, J.A. de Haseth, L.V. Azarraga, "Capillary GC/FTIR", ANAL. CHEM. **55** (1983) 1361A–1387A.
- [91] W. Herres, "HRGC-FTIR: Capillary gas chromatography–fourier transform infrared spectroscopy. Theory and application", Chromatographic Methods Series, Hüthig Verlag, Heidelberg (1987).

- [92] R. White, "Chromatography/fourier transform infrared spectroscopy and its applications", Practical Spectroscopy Series 10, Marcel Dekker, New York (1990).
- [93] A.H. Mohamed, J.A. Caruso, "Element-selective plasma emission detectors for gas chromatography", in ADVANCES IN CHROMATOGRAPHY **26** (1987) 191–227.
- [94] S.J. Hill, M.J. Bloxham, P.J. Worsfold, "Chromatography coupled with inductively coupled plasma atomic emission spectrometry and inductively coupled plasma mass spectrometry. A review", *J. ANAL. AT. SPECTROM.* **8** (4) (1993) 499–515 (CA 119:39674y).
- [95] P.C. Uden, "Element-specific chromatographic detection by atomic absorbtion, plasma atomic emission and plasma mass spectrometry", *J. CHROMATOGR. A* **703** (1995) 393–416.
- [96] S. Pedersen-Bjergaard, T. Greibrokk, "Environmental application of capillary gas chromatography coupled with atomic emission detection. A review", *J. HIGH RESOLUT. CHROMATOGR.* **19** (1996) 597–607.
- [97] R. Buffington, "GC-atomic emission spectroscopy using microwave plasmas", Hewlett-Packard Co.
- [98] K. Sutton, R.M.C. Sutton, J.A. Caruso, "Inductively coupled plasma mass spectrometric detection for chromatography and capillary electrophoresis. Review", *J. CHROMATOGR. A* **789** (1997) 85–126.
- [99] J. Marshall, S. Chinery, E.H. Evans, A. Fisher, "Atomic emission spectrometry", *J. ANAL. AT. SPECTROM.* **13** (1998) R107–R128.
- [100] C.L. Wilkins, G.N. Giss, G.M. Brissey, S. Steiner, "Direct-linked gas chromatography–Fourier transform infrared–mass spectrometer system", *ANAL. CHEM.* **53** (1981) 113–117.
- [101] R.W. Crawford, T. Hirschfeld, R.H. Sanborn, C.M. Wong, "Organic analysis with a combined capillary gas chromatography/mass spectrometer/Fourier transform infrared spectrometer", *ANAL. CHEM.* **54** (1982) 817–820.
- [102] C.L. Wilkins, G.N. Giss, R.L. White, G.M. Brissey, E.C. Onyiriuka, "Mixture analysis by gas chromatography/Fourier transform infrared spectrometry/mass spectrometry", *ANAL. CHEM.* **54** (1982) 2260–2264.
- [103] D.A. Laude, Jr., G.M. Brissey, C.F. Ijames, R.S. Brown, C.L. Wilkins, "Linked gas chromatography/Fourier transform infrared/Fourier transform mass spectrometry with integrated electron impact and chemical ionization", *ANAL. CHEM.* **56** (1984) 1163–1168.
- [104] D.F. Gurka, R. Titus, "Rapid nontarget screening of environmental extracts by direct linked gas chromatography/Fourier transform infrared/mass spectrometry", *ANAL. CHEM.* **58** (1986) 2189–2194.
- [105] C.L. Wilkins, *ANAL. CHEM.* **59** (1987) 571A–581A.
- [106] J.R. Cooper, C.L. Wilkins, "Utilization of spectrometric information in linked gas chromatography–Fourier transform infrared spectroscopy–mass spectrometry", *ANAL. CHEM.* **61** (1989) 1571–1577.
- [107] D.F. Gurka, I. Farnham, B.B. Potter, S. Pyle, R. Titus, W. Duncan, "Quantitation capability of a directly linked gas chromatography/Fourier transform infrared/mass spectrometry system", *ANAL. CHEM.* **61** (1989) 1584–1589.
- [108] W.F. Duncan, P.C. Uden, L.D. Hoey, "CGC/IR/MS analysis of pyrolysis products and additives for polysiloxanes and polybutadienes", in PROC. 10TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 22–25, 1989), P. Sandra, G. Redant, Eds., Hüthig Verlag, Heidelberg (1989) 1219–1229.
- [109] D.T. Williams, Q. Tran, P. Fellin, K.A. Brice, "Evaluation of gas chromatography–Fourier transform infrared spectroscopy–mass spectrometry for the analysis of phenolic compounds", *J. CHROMATOGR.* **549** (1991) 297–311.
- [110] R. Oguchi, A. Shimizu, S. Yamashita, K. Yamaguchi, P. Wylie, "Polymer analysis using pyrolysis-GC-FTIR-MS and GC-AED", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 412–416.
- [111] D.B. Hooker, J. Dezwaan, "Analytical problem solving with simultaneous atomic emission–mass spectrometric detection for gas chromatography", in ACS SYMP. SER. 479 (ELEM.–SPECIFIC CHROMATOGR. DETECT. AT. EMISS. SPECTROSC.) (1992) 132–135 (CA 116:120076q).
- [112] S. Sommer, R. Kamps, S. Schumm, K.F. Kleinermanns, "GC/FTIR/MS spectroscopy of native polychlorinated dibenzo-p-dioxins and dibenzofurans extracted from municipal fly-ash", *ANAL. CHEM.* **69** (1997) 1113–1118.
- [114] K.A. Krock, N. Ragunathan, C.L. Wilkins, "Parallel cryogenic trapping multidimensional gas chromatography with directly linked infrared and mass spectral detection", *J. CHROMATOGR.* **645** (1993) 153–159.

- [115] K.A. Krock, C.L. Wilkins, "Environmental application of combined multidimensional gas chromatography–infrared–mass spectrometry", *ANAL. CHIM. ACTA* **277** (1993) 381–388.
- [116] K.A. Krock, N. Ragunathan, C. Klawun, T. Sasaki, C.L. Wilkins, "Multidimensional gas chromatography–infrared spectrometry–mass spectrometry. Plenary lecture", *ANALYST* **119** (4) (1994) 483–489.
- [117] K.A. Krock, C.L. Wilkins, "Quantitative aspects of a valve-based, multi stage multidimensional gas chromatography–infrared spectroscopy–mass spectrometry system", *J. CHROMATOGR. A.* **678** (1994) 265–277.
- [118] K.A. Krock, N. Ragunathan, C.L. Wilkins, "Multidimensional gas chromatography coupled with infrared and mass spectrometry for analysis of eucalyptus essential oils", *ANAL. CHEM.* **66** (1994) 425–430.
- [119] C.L. Wilkins, "Multidimensional GC for qualitative IR and MS of mixtures", *ANAL. CHEM.* **66** (1994) A295–A301.
- [120] D. Rosenthal, "Theoretical limitations of gas chromatographic/mass spectrometric identification of multicomponent mixtures", *ANAL. CHEM.* **54** (1982) 63–66.
- [121] J. Enqvist, M. Enqvist, "Comparison of two-stage retention index monitoring capillary gas chromatography and selected ion monitoring capillary gas chromatography–mass spectrometry as analytical tools", *J. HIGH RESOLUT. CHROMATOGR.* **17** (1994) 141–144.
- [122] J. Schuberth, "Joint use of retention index and mass spectrum in postmortem tests for volatile organics by headspace capillary gas chromatography with ion-trap detection", *J. CHROMATOGR. A.* **674** (1994) 63–71.
- [123] B.W. Cook, M. Cleary, "Systematic approach to coupled gas chromatography Fourier transform infrared spectroscopy using two wide bore capillary columns", *ANALYST* **113** (1988) 1171–1174.
- [124] C. Bicchi, A. D'Amato, M. Orlandini, "Identification of the constituents of a complex mixture by combined use of retention indices and specific multidetection responses", *J. HIGH RESOLUT. CHROMATOGR.* **17** (1994) 335–338.
- [125] J. Winskowski, "Gas chromatographic identification of compounds from retention indices and by different detectors", *CHROMATOGRAPHIA* **17** (1983) 160.
- [126] N.R. Smyrl, D.M. Hembree, Jr., W.E. Davis, D.M. Williams, J.C. Vance, "Simultaneous GC-FTIR/GC-MS analysis for isomer-specific identification and quantitation of complex mixture compounds", *APPL. SPECTROSC.* **46** (1992) 277–282.
- [127] L. Yongzhi, C. Zhangping, W. Xuanjie, "Allitridin analysis by FID/FPD dual channel gas chromatography", *J. CHROMATOGR. SCI.* **32** (1994) 61–64.
- [128] D.M. Hembree, N.R. Smyrl, W.E. Davis, D. Williams, *ANALYST* **118** (1993) 249.
- [129] P. Karpe, S. Kirchner, P. Rouxel, "Thermal desorption–gas chromatography–mass spectrometry–flame ionization detection–sniffer multi-coupling: a device for the determination of odorous volatile organic compounds in air", *J. CHROMATOGR. A.* **708** (1995) 105–114.
- [130] W. Bertsch, E. Anderson, G. Holzer, "Two-dimensional high resolution GLC environmental analysis, preliminary results", *CHROMATOGRAPHIA* **10** (1977) 449–454.
- [131] R.D. Cox, R.F. Earp, "Determination of trace level organics in ambient air by high-resolution gas chromatography with simultaneous photoionization and flame ionization detection", *ANAL. CHEM.* **54** (1982) 2265–2270.
- [132] H.-J. Stan, D. Mrowetz, "Residue analyzes of pesticides in food by two-dimensional gas chromatography with capillary columns and parallel detection with flame-photometric and electron-capture detection", *J. CHROMATOGR.* **279** (1983) 173–187.
- [133] H.-J. Stan, D. Mrowetz, "Residue analysis of pesticides in food with two-dimensional gas chromatography using capillary columns and parallel detection with FPD and ECD", in *PROC. 5TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, April 26–28, 1983), J. Rijks, Ed., Elsevier, Amsterdam (1983) 189–202.
- [134] A. D'Amato, C. Bicchi, M. Galli, "The use of three way press-fit connectors (Y connectors) in the analysis of essential oils", *J. HIGH RESOLUT. CHROMATOGR.* **12** (1989) 349–352.
- [135] J.R. Hancock, G.R. Peters, "Retention index monitoring of compounds of chemical defence interest using thermal desorption gas chromatography", *J. CHROMATOGR.* **538** (1991) 249–257.
- [136] Z. Penton, "Determination of residual solvent in pharmaceutical preparations by static headspace GC", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 329–331.

- [137] H. Kessels, W. Hoogerwerf, J. Lips, "The determination of volatile organic compounds from EPA method 524.2 using purge-and-trap capillary gas chromatography, ECD, and FID", *J. CHROMATOGR. SCI.* **30** (1992) 247–255.
- [138] C. Bicchi, A. D'Amato, M. Orlandini, "Identification of the constituents of a complex mixture by combined use of retention indices and specific multidetection responses", *J. HIGH RESOLUT. CHROMATOGR.* **17** (1994) 335–338.
- [139] M. Hrvnac, W. Frischknecht, L. Cechova, "Gas chromatographic multidetector coupled to a glass capillary column", *ANAL. CHEM.* **48** (1976) 937–940.
- [140] T.H. Parliment, M.D. Spencer, "Application of simultaneous FID/NPD/FPD detectors in the capillary gas chromatographic analysis of flavors", *J. CHROMATOGR. SCI.* **19** (1981) 435–438.
- [141] C.L. Wilkins, G.N. Giss, R.L. White, G.M. Brissey, E.C. Onyiriuka, "Mixture analysis by gas chromatography/Fourier transform infrared spectrometry/mass spectrometry", *ANAL. CHEM.* **54** (1982) 2260–2264.
- [142] P.T. Williams, K.D. Bartle, D.G. Mills, G.E. Andrews, "Capillary GC analysis of diesel fuels using simultaneous parallel triple detection", *J. HIGH RESOLUT. CHROMATOGR.* **9** (1986) 39–43.
- [143] F. Lindqvist, "Sulfur-specific detection in air by photoionization in a multiple detector gas chromatographic system", *J. HIGH RESOLUT. CHROMATOGR.* **12** (1989) 628–631.
- [144] K.K. Gaines, W.H. Chatham, S.O. Farwell, "Comparison of the SCD and FPD for HRGC determination of atmospheric sulfur gases", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 489–493.
- [145] H.-J. Stan, S. Heil, "Two-dimensional capillary gas chromatography with three selective detectors as a valuable tool in residue analysis. State-of-the art", *FRESENIUS' J. ANAL. CHEM.* **339** (1991) 34–39.
- [146] H.-J. Stan, B. Christall, "Residue analysis of onions and other foodstuffs with a complex matrix using two-dimensional capillary-GC with three selective detectors", *FRESENIUS' J. ANAL. CHEM.* **339** (1991) 395–398.
- [147] R.J. Bohler, A.J. McCormack, J.M. McCann, "Simultaneous detection of aromatics, sulfur and hydrocarbons in diesel fuels by gas chromatography", *PREP.–AM. CHEM. SOC., DIV. PET. CHEM.* **36** (2) (1991) 204–207 (CA 115:186487k).
- [148] F. Bruner, P. Ciccioli, S. Zelli, "Improved double detection gas chromatograph–mass spectrometer interface for the analysis of complex organic mixtures", *ANAL. CHEM.* **45** (1973) 1002–1006.
- [149] Y. Hoshika, G. Muto, "Capillary gas chromatography of lower aliphatic C₂–C₅ aldehydes in the free form as a group of pungent odorants in air", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 330–334.
- [150] H.A. McLeod, A.G. Butterfield, D. Lewis, W.E.J. Phillips, D.E. Coffin, "Gas-liquid chromatography system with flame ionization, phosphorus, sulfur, nitrogen, and electron capture detectors operating simultaneously for pesticide residue analysis", *ANAL. CHEM.* **47** (1975) 674–679.
- [151] C.L. Wilkins, G.N. Giss, R.L. White, G.M. Brissey, E.C. Onyiriuka, "Mixture analysis by gas chromatography/Fourier transform infrared spectrometry/mass spectrometry", *ANAL. CHEM.* **54** (1982) 2260–2264.
- [152] W. Herres, H. Idstein, P. Schreier, "Flavor analysis by HRGC/FT-IR: cherimoya (*Annona Cherimolia*, mill.) fruit volatiles", *J. HIGH RESOLUT. CHROMATOGR.* **6** (1983) 590–594.
- [153] R. Deleu, A. Copin, "Multidetection of pesticides in capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **7** (1984) 338–339.
- [154] W.F. Duncan, P.C. Uden, L.D. Hoey, "CGC/IR/MS analysis of pyrolysis products and additives for polysiloxanes and polybutadienes", in *PROC. 10TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 22–25, 1989), P. Sandra, G. Redant, Eds., Hüthig Verlag, Heidelberg (1989) 1219–1229.
- [155] N. Kirshen, E. Almasi, "The automated determination of volatile organic compounds in ambient air and/or soil gas by gas chromatography with selective detectors", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 484–489.
- [156] F.L. Herman, "Tandem detectors to quantitative overlapping chromatographic peaks", *ANAL. CHEM.* **65** (1993) 1023–1027.
- [157] G.R. Verga, F. Pigozzo, "Simultaneous selective determination of organic pollutants by dual detection system", in *PROC. 18TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 20–24, 1996), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1996) 1532–1536.

- [158] G.P. Mapelli, G.R. Verga, F. Pigozzo, "Head space determination of volatile compounds in complex matrices by using simultaneous selective detectors", in PROC. 18TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 20-24, 1996), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1996) 1628–1632.
- [159] S. Sommer, R. Kamps, S. Schumm, K.F. Kleinermanns, "GC/FT-IR/MS spectroscopy of native polychlorinated dibenzo-p-dioxins and dibenzofurans extracted from municipal fly-ash", ANAL. CHEM. **69** (1997) 1113–1118.
- [160] H.J. Schroers, E. Jermann, J. Begerow, H. Hajimiragh, A.M. Chiarottimar, L. Dunemann, "Determination of physiological levels of volatile organic compounds in blood using static headspace capillary gas chromatography with serial triple detection", ANALYST **123** (1998) 715–720.
- [161] Y.C. Chen, J.G. Lo, "Gas chromatography with flame ionization and flameless sulfur chemiluminescence detectors in series for dual channel detection of sulfur compounds", CHROMATOGRAPHIA **43** (1996) 522–526.

Hivatkozások a 2.3. szakaszban

- [1] A.T. James, A.J.P. Martin, "Gas-liquid partition chromatography: the separation and microestimation of volatile fatty acids from formic acid to dodecanoic acid", BIOCHEM. J. **50** (1952) 679–690 (CA 46:4884a).
- [2] A.T. James, A.J.P. Martin, "Gas-liquid partition chromatography. a technique of the analysis of volatile minerals", ANALYST **77** (1952) 915–932 (CA 47:1529d).
- [3] A.T. James, A.J.P. Martin, G.H. Smith, "Gas-liquid partition chromatography. separation and microestimation of ammonia and the methylamines", BIOCHEM. J. **52** (1952) 238–242 (CA 47:446d).
- [4] B.W. Bradford, D. Harvey, D.E. Chalkley, "The chromatographic analysis of hydrocarbon mixtures", J. INST. PETROLEUM **41** (1955) 80–91 (CA 49:7839h).
- [5] A.C. Cope, N.A. Lebel, H.-H. Lee, W.R. Moore, "Amine oxides. Part III. Selective formation of olefins from unsymmetrical amine oxides and quaternary ammonium hydroxides", J. AM. CHEM. SOC. **79** (1957) 4720–4729 (CA 52:2769h).
- [6] A.C. Cope, C.L. Bumgardner, E.E. Schweizer, "Amine oxides. Part IV. alicyclic olefins from amine oxides and quaternary ammonium hydroxides", J. AM. CHEM. SOC. **79** (1957) 4729–4733 (CA 52:2771c).
- [7] E. Gil-Av, J. Herling, J. Shabtai, "Gas-liquid partition chromatography of mixtures of the three isomeric methylcyclohexenes and methylcyclohexane", CHEM. IND. (London) (1957) 1483–1484 (CA 52:4413i).
- [8] M.E. Bednas, D.S. Russel, "Silver nitrate solutions in gas chromatography", CAN. J. CHEM. **36** (1958) 1272–1276 (CA 53:1979a).
- [9] A.C. Cope, E.M. Acton, "Amine oxides. Part V. Olefins from N,N-dimethylmenthylamine and N,N-dimethylneomenthylamine oxides", J. AM. CHEM. SOC. **80** (1958) 355–359 (CA 52:10006g).
- [10] A.D. Zorin, A.E. Ezheleva, G.G. Devyatikh, "Analysis of butane-butylene fractions by gas-liquid partition chromatography", TRUDY KHIM. I KHIM. TEKHNOL. **1** (1958) 605–610 (CA 54: 6385g).
- [11] H.M. Tenney, "Selectivity of various liquid substrates used in gas chromatography", ANAL. CHEM. **30** (1958) 2–8.
- [12] J.A. Barnard, H.W.D. Hughes, "Analysis of light hydrocarbon gas mixtures", NATURE **183** (1959) 250.
- [13] J.W. Powell, M.C. Whiting, "Decomposition of sulfonylhydrazone salts. Part I. Mechanism and stereochemistry", TETRAHEDRON **7** (1959) 305–310 (CA 54:7619g).
- [14] J. Shabtai, J. Herling, E. Gil-Av, "Gas-liquid partition chromatography of isomeric alkylcyclopentenes and alkylidenecyclopentanes", J. CHROMATOGR. **2** (1959) 406–410 (CA 54:4269f).
- [15] F. Armitage, "Analysis of isoprene by gas chromatography", J. CHROMATOGR. **2** (1959) 655–657 (CA 54:20672i).
- [16] B. Smith, R. Ohlson, "Gas chromatographic separation of 3- and 4-methyl-1-pentene", ACTA CHEM. SCAND. **13** (1959) 1253–1253 (CA 55:25725e).
- [17] A.C. Cope, P.T. Moore, W.R. Moore, "Equilibration of cis- and trans-cycloalkenes", J. AM. CHEM. SOC. **82** (1960) 1744–1749 (CA 54:14886a).

- [18] R.A. Hively, "Identification of hydrocarbons by gas chromatography", *J. CHEM. ENG. DATA* **5** (1960) 237–240 (CA 54:20668h).
- [19] J.W. Powell, M.C. Whiting, "Analysis of mixtures of octahydronaphthalenes", *TETRAHEDRON* **12** (1961) 163–167 (CA 55:16494d).
- [20] E. Bendel, M. Kern, R. Janssen, G. Steffan, "Separation by gas chromatography of all double bond isomers of octene", *ANGEW. CHEM.* **74** (1962) 905–906 (CA 58:2827c).
- [21] J. Herling, J. Shabtai, E. Gil-Av, "Gas chromatography with stationary phases containing silver nitrate. Part III. Isomeric C₈ and C₉ cyclohexenes and p-menthenes", *J. CHROMATOGR.* **8** (1962) 349–354 (CA 58:7375f).
- [22] B. Smith, R. Ohlson, "Gas chromatographic separation of unsaturated hydrocarbons using silver nitrate in ethylene glycol as the stationary phase", *ACTA CHEM. SCAND.* **16** (1962) 351–358.
- [23] J. Shabtai, J. Herling, E. Gil-Av, "Gas chromatography with stationary phases containing silver nitrate. Part IV. The methylcyclobutenes, methylenecyclobutane, C₅ dienes, and related compounds", *J. CHROMATOGR.* **11** (1963) 32–37 (CA 59:9292g).
- [24] R.J. Cvetanovic, F.J. Duncan, W.E. Falconer, "Gas chromatographic separation of deuterium-substituted olefins by unstable complex formation", *CAN. J. CHEM.* **41** (1963) 2095–2097 (CA 59: 9292d).
- [25] E.K.C. Lee, F.S. Rowland, "Separation of isotopic position isomers of tritium-labeled olefins by gas chromatography", *ANAL. CHEM.* **36** (1964) 2181–2183.
- [26] A. Zlatkis, G.S. Chao, H.R. Kaufman, "Substrates containing silver nitrate for gas chromatography", *ANAL. CHEM.* **36** (1964) 2354.
- [27] E. Gil-Av, J. Herling, "Complex forming stationary phases in gas liquid partition chromatography", in *PROC. CHROMATOGR. METHODS IMMED. SEPAR.* (Athens, 1965) **1** (1966) 167–176 (CA 68:38625z).
- [28] J. Shabtai, "Gas chromatography with stationary phases containing silver-nitrate. Part VI. Methylcycloheptenes and methylenecycloheptane", *J. CHROMATOGR.* **18** (1965) 302–307 (CA 63: 10650f).
- [29] J.V. Mortimer, P.L. Gent, "Use of modified Bentone-34 for the gas chromatographic separation of aromatic hydrocarbons", *NATURE* **197** (1963) 789–790.
- [30] A.F. Spencer, "Rapid separation of xylenes and ethylbenzene by gas chromatography using packed columns", *ANAL. CHEM.* **35** (1963) 592.
- [31] M. van der Stricht, J. van Rysselberge, "Modified Bentone 34, a new addition to stationary phases in gas chromatography to assist in separation of difficult aromatic mixtures", *J. GAS CHROMATOGR.* **1** (8) (1963) 29–33 (CA 59:10746b).
- [32] J.V. Mortimer, P.L. Gent, "The use organo-clays as gas chromatographic stationary phases", *ANAL. CHEM.* **36** (1964) 754–756.
- [33] E.W. Cieplinski, "Gas chromatographic analysis of aromatic hydrocarbons with modified bentonite columns. Effect of Bentone-34 concentration on performance of packed columns", *ANAL. CHEM.* **37** (1965) 1160–1161.
- [34] J.H. Jordan, "Analysis of hydrocarbons using mixed 1,2,3-tris(2-cyanoethoxy)propane and SE-30 gum rubber liquid phase", *J. GAS CHROMATOGR.* **2** (1964) 346.
- [35] A. Zlatkis, S.Y. Ling, H.R. Kaufman, "Resolution of isomeric xylenes by gas-liquid chromatography", *ANAL. CHEM.* **31** (1959) 945–947.
- [36] A.I.M. Keulemans, A. Kwanten, P. Zaal, "The selectivity of the stationary liquid in vapour phase chromatography", *ANAL. CHIM. ACTA* **13** (1955) 357–372.
- [37] H. Miyake, M. Mitooka, "Gas chromatographic analysis of C₁-C₄ hydrocarbons by using mixed stationary liquid of high polarity", *NIPPON KAGAKU ZASSHI* **84** (1963) 923–928 (CA 61:7677d).
- [38] H. Miyake, M. Mitooka, "Gas chromatographic analysis of C₁-C₄ hydrocarbons with mixed stationary liquid of Apiezon–dibutyl maleate mixtures", *NIPPON KAGAKU ZASSHI* **84** (1963) 593–595 (CA 61:1275g).
- [39] A. Zlatkis, "Resolution of isomeric hexanes by gas-liquid chromatography", *ANAL. CHEM.* **30** (1958) 332–333.
- [40] J.Q. Walker, D.L. Ahlberg, "Quantitative analysis of aromatic hydrocarbons by capillary gas chromatography", *ANAL. CHEM.* **35** (1963) 2022–2027.
- [41] R.D. Schwartz, D.J. Brasseaux, "Resolution of complex hydrocarbon mixtures by capillary column gas liquid chromatography", *ANAL. CHEM.* **35** (1963) 1374–1382.

- [42] H. Miyake, M. Mitooka, "Gas chromatographic analysis of gaseous hydrocarbons with a capillary column", NIPPON KAGAKU ZASSHI **85** (1964) 326–331 (CA 61:12604c).
- [43] H. Miyake, M. Mitooka, T. Matsumoto, "Studies of gas chromatography with mixed stationary liquids. Part IV. The analysis of aromatic hydrocarbons from C₆ to C₁₀ with a capillary column", BUL. CHEM. SOC. JAPAN **38** (1965) 1062–1071.
- [44] W.H. McFadden, "Use of mixed stationary liquids in gas-liquid chromatography", ANAL. CHEM. **30** (1958) 479–481.
- [45] P.P. Nair, C. Bucana, S. de Leon, D.A. Turner, "Gas chromatographic studies of vitamins D₂ and D₃", ANAL. CHEM. **37** (1965) 631–636.
- [46] R.L. Gatrell, "A mixed-substrate column for gas chromatographic analysis of lacquer thinners", ANAL. CHEM. **35** (1963) 923–924.
- [47] A.H. Duvall, W.F. Tully, "Gas-liquid chromatography of mixtures containing phenol and five of its tert-butyl derivatives", J. CHROMATOGR. **11** (1963) 38–44.
- [48] K.D. Parker, C.R. Fontan, P.L. Kirk, "Improved gas chromatographic column for barbiturates", ANAL. CHEM. **35** (1963) 418–419.
- [49] K.A. McCully, W.P. McKinley, "Determination of chlorinated pesticide residues in fat by electron capture gas chromatography", J. ASSOC. OFFIC. AGR. CHEMISTS **47** (4) (1964) 652–658 (CA 61:11238e).
- [50] F. Ascoli, B. Pispisa, F. Servello, "The use of pentabenzoyl- α -glucose as stationary phase in gas chromatography", J. CHROMATOGR. **6** (1961) 544–546.
- [51] E. Gil-Av, J. Herling, "Determination of the stability constants of complexes by gas chromatography", J. PHYS. CHEM. **66** (1962) 1208–1209 (CA 58:64g).
- [52] M.A. Muhs, F.T. Weiss, "Determination of equilibrium constants of silver-olefin complexes using gas chromatography", J. AM. CHEM. SOC. **84** (1962) 4697.
- [53] G.F. Freeguard, R. Stock, "Some static measurement on gas-liquid chromatographic systems involving dinonyl phthalate and squalane", in GAS CHROMATOGRAPHY (1962), M. van Swaay, Ed., Butterworths, London (1962) 102–110.
- [54] R.J. Cvetanovic, F.J. Duncan, W.E. Falconer, R.S. Irwin, "Secondary deuterium isotope effects on the stability of the silver ion-olefin complexes", J. AM. CHEM. SOC. **87** (1965) 1827.
- [55] E. Gil-Av, J. Herling, "Complex forming stationary phases in gas liquid partition chromatography", in PROC. CHROMATOGR. METHODS IMMED. SEPAR. (Athens, 1965) **1** (1966) 167–176 (CA 68:38625z).
- [56] A.I.M. Keulemans, A. Kwantes, P. Zaal, "The selectivity of the stationary liquid in vapour phase chromatography", ANAL. CHIM. ACTA **13** (1955) 357–372.
- [57] F. van de Craats, "Application of vapour phase chromatography in the gas-analytical field", ANAL. CHIM. ACTA **14** (1956) 136–149.
- [58] E.M. Fredericks, F.R. Brooks, "Analysis of gaseous hydrocarbons by gas-liquid partition chromatography", ANAL. CHEM. **28** (1956) 297–303.
- [59] M.E. Bednas, D.S. Russel, "Silver nitrate solutions in gas chromatography", CAN. J. CHEM. **36** (1958) 1272–1276 (CA 53:1979a).
- [60] J.A. Barnard, H.W.D. Hughes, "Analysis of light hydrocarbon gas mixtures", NATURE **183** (1959) 250.
- [61] G.R. Primavesi, "Analysis of light hydrocarbons", NATURE **184** (1959) 2010–2011.
- [62] W.H. McFadden, "Use of mixed stationary liquids in gas-liquid chromatography", ANAL. CHEM. **30** (1958) 479–481.
- [63] A.D. Zorin, A.E. Ezheleva, G.G. Devyatlykh, "Analysis of butane-butylene fractions by gas-liquid partition chromatography", TRUDY KHIM. I KHIM. TEKHNOL. **1** (1958) 605–610 (CA 54: 6385g).
- [64] A. Zlatkis, "Resolution of isomeric hexanes by gas-liquid chromatography", ANAL. CHEM. **30** (1958) 332–333.
- [65] L. Rohrschneider, "Polarity of the stationary phase in gas chromatography", Z. ANAL. CHEM. **170** (1959) 256–263 (CA 54:3045i).
- [66] H.J. Maier, O.C. Karpathy, "Prediction of separation and specifications of chromatographic columns", J. CHROMATOGR. **8** (1962) 308–318.
- [67] G.P. Hildebrand, C.N. Reilley, "Use of combination columns in gas liquid chromatography", ANAL. CHEM. **36** (1964) 47–58.

- [68] M. Singliar, A. Bobak, J. Brida, L. Lukacovic, "Mixed stationary phases for gas-liquid chromatography", *Z. ANAL. CHEM.* **177** (1960) 161–166 (CA 55:19614b).
- [69] A.A. Zhukhovitskii, M.S. Selenkina, N.M. Turkel'taub, "Consecutive columns in gas chromatography", *ZH. FIZ. KHM.* **36** (1962) 993–998 (CA 57:6581i).
- [70] R.A. Keller, G.H. Stewart, "Mixed solvents in gas liquid chromatography", *ANAL. CHEM.* **36** (1964) 1186–1191.
- [71] A. Waksmundzki, E. Soczewinski, Z. Suprynowicz, "The relation between the composition of the mixed stationary phase and the retention time in gas-liquid partition chromatography", *COLLECTION CZECH. CHEM. COMMUN.* **27** (1962) 2001–2006 (CA 58:11977f).
- [72] A. Waksmundzki, Z. Suprynowicz, "Separation of methyl esters of fatty acids by gas-liquid partition chromatography using a binary liquid phase", *CHEM. ANAL. (Warsaw)* **10** (3) (1965) 377–385 (CA 64:9965c).
- [73] A. Waksmundzki, Z. Suprynowicz, "The use of binary stationary phases in gas-liquid partition chromatography", *J. CHROMATOGR.* **18** (1965) 232–242.
- [74] A. Waksmundzki, Z. Suprynowicz, T. Pietrusinska, "Application of binary stationary phases for separation of hexane isomers (petroleum ether) by partition gas chromatography", *CHEM. ANAL. (Warsaw)* **10** (3) (1965) 367–375 (CA 64:12531a).
- [75] A. Waksmundzki, Z. Suprynowicz, I. Miedziak, "Combination columns in partition gas chromatography", *CHEM. ANAL. (Warsaw)* **12** (6) (1967) 1205–1211 (CA 68:119000u).
- [76] A. Waksmundzki, Z. Suprynowicz, I. Miedziak, "Effect of amount of binary mixed phase on the adsorption properties of support in gas chromatography", *J. GAS CHROMATOGR.* **4** (2) (1966) 74–76 (CA 64:11899e).
- [77] T. Czajkowska, A. Waksmundzki, "The properties of binary stationary phases in gas-liquid partition chromatography. Complex association systems", *J. CHROMATOGR.* **119** (1976) 91–98.
- [78] T. Czajkowska, "Properties of binary stationary phases in gas-liquid partition chromatography. Part II. Influence of the amount and polarity of stationary phase on the observed retention data", *J. CHROMATOGR.* **241** (1982) 9–18.
- [79] T. Czajkowska, "Properties of binary stationary phases in gas-liquid partition chromatography. Part III. Application of association constants for the determination of the adsorptive properties of the support", *J. CHROMATOGR.* **243** (1982) 245–252 (CA 97:79459r).
- [80] A. Waksmundzki, Z. Suprynowicz, I. Miedziak, "Use of binary liquid phases for the identification of organic substances on triangular diagrams in gas chromatography", *CHEM. ANAL. (Warsaw)* **13** (3) (1968) 635–640 (CA 70:16878y).
- [81] J.C. Giddings, "Plate height of nonuniform chromatographic columns. Gas compression effects, coupled columns, and analogous systems", *ANAL. CHEM.* **35** (1963) 353–356.
- [82] J. Kwok, L.R. Snyder, J.C. Sternberg, "Efficiency of chromatographic columns connected in series. Nonadditivity of theoretical plates in chromatography", *ANAL. CHEM.* **40** (1968) 118–122.
- [83] R.S. Porter, R.L. Hinkins, L. Tornheim, J.F. Johnson, "Computer optimization of mixed liquid phases for gas chromatography", *ANAL. CHEM.* **36** (1964) 260–262.
- [84] A.B. Littlewood, F.W. Willmott, "Experimental and theoretical study of specific retention volumes in mixtures of polar and nonpolar stationary liquids", *ANAL. CHEM.* **38** (1966) 1031–1041.
- [85] A.B. Littlewood, F.W. Willmott, "The identity of equivalent mixtures as stationary liquids", *J. GAS CHROMATOGR.* **5** (1967) 543–545.
- [86] J.B. Carmichael, "Theoretical consideration regarding tandem column arrangements with application to gas-liquid and gel permeation chromatography", *SEPAR. SCI.* **3** (3) (1968) 249–254, (CA 70: 61406d).
- [87] C.L. Young, "Mixed solvents in gas-liquid chromatography", *J. CHROMATOGR. SCI.* **8** (1970) 103–104.
- [88] M.S. Vigdergauz, M.I. Afanas'ev, "Separation of substances on mixed sorbents", *GAZOV. KROMATOGR.* **4** (1966) 24–28 (CA 68:6636n).
- [89] J.F.K. Huber, E. Kenndler, H. Markens, "Adjustment of stationary phase selectivity in gas-liquid chromatography by using mixed liquid phases", *J. CHROMATOGR.* **167** (1978) 291–305.

Hivatkozások a 2.4. szakaszban

- [1] G.J. Price, "The use and properties of mixed stationary phases in gas chromatography", in ADVANCES IN CHROMATOGRAPHY, J.C. Giddings, E. Grushka, P.R. Brown, Eds., **28** (1989) 113–163.
- [2] J.H. Purnell, J.M. Vargas de Andrade, "Solution and complexing studies. Part I. Gas-liquid chromatographic investigation of supposed complexing systems", *J. AM. CHEM. SOC.* **97** (1975) 3585–3590.
- [3] J.H. Purnell, J.M. Vargas de Andrade, "Solution and complexing studies. Part II. Comparison of nuclear magnetic resonance and gas-liquid chromatographic data", *J. AM. CHEM. SOC.* **97** (1975) 3590–3593.
- [4] R.J. Laub, J.H. Purnell, "Solution and complexing studies. Part III. Evidence for a microscopic partitioning theory of solution", *J. AM. CHEM. SOC.* **98** (1976) 30–35.
- [5] R.J. Laub, J.H. Purnell, "Solution and complexing studies. Part IV. Extension of microscopic partition theory to include nonstoichiometric complexation and solvent effects, and the correlation of GLC, UV, and NMR data", *J. AM. CHEM. SOC.* **98** (1976) 35–39.
- [6] R.J. Laub, J.H. Purnell, "Criteria for the use of mixed solvents in gas-liquid chromatography", *J. CHROMATOGR.* **112** (1975) 71–79.
- [7] R.J. Laub, J.H. Purnell, "Quantitative approach to the use of multicomponent solvents in gas-liquid chromatography", *ANAL. CHEM.* **48** (1976) 799–803.
- [8] R.J. Laub, J.H. Purnell, "Optimization of gas chromatographic analysis of complex mixtures of unknown composition", *ANAL. CHEM.* **48** (1976) 1720–1724.
- [9] W.K. Al-Thamir, R.J. Laub, J.H. Purnell, "Gas chromatographic separation of all C₁–C₄ hydrocarbons by multi-substrate gas-solid-liquid chromatography", *J. CHROMATOGR.* **142** (1977) 3–14.
- [10] R.J. Laub, J.H. Purnell, D.M. Summers, P.S. Williams, "Employment of relative retention times and capacity factors for the prediction of optimized gas-liquid chromatographic separations", *J. CHROMATOGR.* **155** (1978) 1–8.
- [11] R.J. Laub, J.H. Purnell, "Chromatographic analysis of mixtures of unknown complexity. Use of relative retention and capacity factor data", *J. CHROMATOGR.* **161** (1978) 59–68.
- [12] J.H. Purnell, "Strategy for chromatography", in *PROC. ANAL. DIV. CHEM. SOC.* **16** (4) (1979) 136–139 (CA 91:133408q).
- [13] C.F. Chien, R.J. Laub, M.M. Kopecni, "Utility of mixed packings in gas-liquid chromatography", *ANAL. CHEM.* **52** (1980) 1402–1407.
- [14] C.F. Chien, R.J. Laub, M.M. Kopecni, "Replication of gas-liquid chromatographic retentions with silicone stationary phases", *ANAL. CHEM.* **52** (1980) 1407–1411.
- [15] R.J. Laub, "Retention index schemes for prediction of chromatographic separations", *ANAL. CHEM.* **52** (1980) 1219–1221.
- [16] M.Y.B. Othman, J.H. Purnell, P. Wainwright, P.S. Williams, "Total optimization of the separation of C₁–C₃ chlorohydrocarbons by gas-liquid chromatography", *J. CHROMATOGR.* **289** (1984) 1–16.
- [17] S.G.A.H. Ali, J.H. Purnell, P.S. Williams, "Window analysis optimization of gas chromatographic separations using mixed Porapaks", *J. CHROMATOGR.* **302** (1984) 119–133.
- [18] C.F. Chien, M.M. Kopecni, R.J. Laub, "Multiple and bonded phases in open-tubular column gas chromatography. Blended dimethyl and methylphenyl polysiloxane solvents", *J. CHROMATOGR. SCI.* **22** (1984) 1–6.
- [19] R.J. Laub, J.H. Purnell, P.S. Williams, "Computer optimization of multi-component sorbents in chromatography", *J. CHROMATOGR.* **95** (1977) 135–143.
- [20] R.J. Laub, J.H. Purnell, P.S. Williams, "Computer-assisted prediction of gas chromatographic separation", *J. CHROMATOGR.* **134** (1977) 249–261.
- [21] P.F. Tiley, "Application of conventional thermodynamic approximations to mixed-solvent gas-liquid chromatography", *J. CHROMATOGR.* **179** (1979) 247–257.
- [22] J.H. Purnell, P.S. Williams, G.A. Zabierek, "Optimization of solvent choice and analysis in GLC", in *PROC. 4TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Bad Duerkheim, FRG, 1981), R.E. Kaiser, Ed., Institute of Chromatography, Bad Duerkheim (1981) 573–599 (CA 98:27057j).
- [23] P.J. Schoenmakers, "Optimization of chromatographic selectivity. A guide to method development", *Journal of chromatography library*, Vol. 35, Elsevier, Amsterdam (1986).

- [24] R.J. Laub, J.H. Purnell, "Quantitative optimization of system variables for chromatographic separations. The column temperature", *J. CHROMATOGR.* **161** (1978) 49–57.
- [25] R.J. Laub, „Solute retention in column liquid chromatography. Part III. Computer optimization of mobile-phase composition: program Window”, *J. LIQUID CHROMATOGR.* **7** (1984) 647–660.
- [26] R.J. Laub, "Recent advances in optimization strategies: chromatography, electrochemistry and spectroscopy", *AM. LAB. (FAIRFIELD CONN.)* **13** (3) (1981) 47–56 (CA 94:145734t).
- [27] R.J. Laub, "Recent advances in optimization strategies: chromatography, electrochemistry, and spectroscopy", *INT. LAB.* **11** (4) (1981) 16–29.
- [28] D.F. Lynch, F.A. Palocsay, J.J. Leary, "An extension of the mixed bed column technique in gas chromatography", *J. CHROMATOGR. SCI.* **13** (1975) 533–534.
- [29] J.R. Mann, S.T. Preston, Jr, "Selection of preferred liquid phases", *J. CHROMATOGR. SCI.* **11** (1973) 216–220.
- [30] J.F. Parcher, J.R. Hansbrough, A.M. Koury, "An investigation of phenylmethylsilicone stationary phases and their use in mixed bed columns", *J. CHROMATOGR. SCI.* **16** (1978) 183–189.
- [31] A.S. Melnikov, B.V. Aivazov, "Binary liquid phase of optimum selectivity in gas-liquid chromatography", *ZH. ANAL. KHM.* **28** (4) (1973) 791–796 (CA 79:38304w).
- [32] A.S. Melnikov, B.V. Aivazov, "Choice of three and four component liquid phases of optimum selectivity in gas liquid chromatography", *ZH. ANAL. KHM.* **29** (2) (1974) 365–371 (CA 81:9494u).
- [33] P.L. Fitzgerald, T.N. Gallaher, F.A. Palocsay, J.J. Leary, "Mixing of liquid phases in gas chromatography columns", *J. CHROMATOGR. SCI.* **15** (1977) 119–120.
- [34] J.A. Garcia Dominguez, E. Fernandez Sanchez, J. Garcia Munoz, M.J. Molera, "Ageing of columns of homogeneous mixed stationary phases in gas chromatography", *J. CHROMATOGR. SCI.* **17** (1979) 281–284.
- [35] M.J. Molera, J.A. Garcia Dominguez, E. Pertierre Rimada, "Gas-liquid chromatography with mixed stationary phase", *AN. QUIM.* **64** (12) (1968) 1105–1110 (CA 71:27278z).
- [36] M.J. Molera, J.A. Garcia Dominguez, J. Fernandez Biarge, "Mixed columns made to order in gas chromatography. Part I. Isothermal analysis", *J. CHROMATOGR. SCI.* **7** (1969) 305–312.
- [37] M.J. Molera, J.A. Garcia Dominguez, J. Fernandez Biarge, "Mixed column made to order in gas chromatography. Part II. Programmed temperature analysis at constant pressure drop", *J. CHROMATOGR. SCI.* **11** (1973) 538–546.
- [38] M.J. Molera, J.A. Garcia Dominguez, J. Fernandez Biarge, "Mixed columns made to order in gas chromatography. Part III. Programmed temperature analysis at constant flow rate", *J. CHROMATOGR. SCI.* **14** (1979) 299–302.
- [39] M.D. Cabezudo, E.F. Gorostiza, M. Herraiz, J. Fernandez Biarge, J.A. Garcia Dominguez, M.J. Molera, "Mixed columns made to order in gas chromatography. Part IV. Isothermal selective separation of alcoholic and acetic fermentation products", *J. CHROMATOGR. SCI.* **16** (1978) 61–67.
- [40] G. Dahlmann, H.J.K. Koeser, H.H. Oelert, "Mixed preferred stationary phases in gas-liquid chromatography", *J. CHROMATOGR. SCI.* **17** (1979) 307–313.
- [41] G. Yang, W. Xu, Y. Peng, W. Dong, "Selection of the optimal ratio of mixed stationary phase column with computer", *GAODENG XUEXIAO HUAXUE XUEBAO* **10** (7) (1989) 691–695 (CA 112: 151024t).
- [42] A. Gröbler, G. Bálizs, "Investigation on mixed gas chromatographic stationary phases. Part I. Dependence of the retention index on the composition of the stationary phase", *J. CHROMATOGR. SCI.* **17** (1979) 631–635.
- [43] A. Gröbler, G. Bálizs, "Investigation on mixed gas chromatographic stationary phases. Part II. Mathematical treatment of the retention index as a function of the stationary phase composition", *J. CHROMATOGR. SCI.* **17** (1979) 671–676.
- [44] A. Gröbler, G. Bálizs, "Investigation on mixed gas chromatographic stationary phases. Part III. A generalized approximation of retention indices for polar-nonpolar stationary phase mixtures", *J. CHROMATOGR. SCI.* **19** (1981) 46–51.
- [45] Gröbler A., "A retenciós indexek változása kevert állófázisokon. IV.", *Magyar Kémikusok Lapja* **40** (1985) 121–125.
- [46] R.L. Pecsok, J. Apffel, "Prediction of gas chromatographic retention behavior with mixed liquid phases", *ANAL. CHEM.* **51** (1979) 594–598.
- [47] W.E. Acree, Jr., J.H. Rytting, "Comment on the prediction of gas chromatographic retention behavior with mixed liquid phases", *ANAL. CHEM.* **52** (1980) 1764–1765.

- [48] M. Lafosse, M. Durand, "Study of squalane–1,3,5-tricyanopentane mixtures by gas-liquid chromatography. Comparison of the dissolution of several solutes in nitrile phases", *J. CHROMATOGR.* **171** (1979) 183–194 (CA 91:9817h).
- [49] M. Lafosse, "Blending bulk dissolution and gas-liquid-adsorption with the mixing of two liquid stationary phases to increase selectivity", *CHROMATOGRAPHIA* **14** (1981) 648–652.
- [50] H. Miyake, M. Mitooka, "Gas chromatographic analysis of C1–C4 hydrocarbons with mixed stationary liquid of Apiezon–dibutyl maleate mixtures", *NIPPON KAGAKU ZASSHI* **84** (1963) 593–595 (CA 61:1275g).
- [51] H. Miyake, M. Mitooka, "Gas chromatographic analysis of C1–C4 hydrocarbons by using mixed stationary liquid of high polarity", *NIPPON KAGAKU ZASSHI* **84** (1963) 923–928 (CA 61:7677d).
- [52] H. Miyake, M. Mitooka, "Gas chromatographic analysis of gaseous hydrocarbons with a capillary column", *NIPPON KAGAKU ZASSHI* **85** (1964) 326–331 (CA 61:12604c).
- [53] H. Miyake, M. Mitooka, T. Matsumoto, "Studies of gas chromatography with mixed stationary liquids. Part IV. The analysis of aromatic hydrocarbons from C₆ to C₁₀ with a capillary column", *BUL. CHEM. SOC. JAPAN* **38** (1965) 1062–1071.
- [54] M. Mitooka, "Chromatography with a mixed stationary phase. Part V. Relation between retention index of oxygenated aliphatic compounds and composition of the stationary phase in mixed-packing gas chromatography", *BUNSEKI KAGAKU* **20** (1971) 1542–1549 (CA 79:41738w).
- [55] M. Mitooka, "Gas chromatography with mixed stationary phase. Part VI. Retention behaviors of aromatic oxygen compounds on gas chromatography with mixed packing of squalane and polar liquids", *BUNSEKI KAGAKU* **21** (1972) 189–196 (CA 77:74640v).
- [56] M. Mitooka, "Gas chromatography with mixed stationary phases. Part VII. Relation between retention indexes of various compounds and relative polarities of stationary phases on mixed packing gas chromatography", *BUNSEKI KAGAKU* **21** (1972) 197–206 (CA 77:9873j).
- [57] M. Mitooka, "Gas chromatography with mixed stationary phase. Part VIII. Effect of the stationary liquid amount and of the property of solid support on retention of solute in mixed packing in gas chromatography", *BUNSEKI KAGAKU* **21** (1972) 354–363 (CA 77:96638c).
- [58] M. Mitooka, "Gas chromatography with mixed stationary phase. Part IX. Consideration of column efficiency and reducing effects of tailing in mixed packing gas chromatography", *BUNSEKI KAGAKU* **21** (1972) 614–622 (CA 77:96637b).
- [59] M. Mitooka, "Gas chromatography with a mixed stationary phase. Part X. Thermodynamic consideration of retention of solute in mixed packing gas chromatography with squalane/polyethylene glycol", *BUNSEKI KAGAKU* **21** (1972) 717–729 (CA 77:119037x).
- [60] M. Mitooka, "Gas chromatography with a mixed stationary phase. Part XI. Consideration of effect of substituent on retention of aromatic oxygen and nitrogen compounds by means of mixed packing gas chromatography", *BUNSEKI KAGAKU* **21** (1972) 729–736 (CA 77:100530f).
- [61] M. Mitooka, "Gas chromatography with a mixed stationary phase. Part XII. Comparison of mixed packing with a mixed stationary liquid in gas chromatography", *BUNSEKI KAGAKU* **21** (1972) 867–877 (CA 77:93166m).
- [62] M. Mitooka, "Gas chromatography with mixed stationary phase. Part XIII. Determination of an optimum composition of stationary phase in mixed packing gas chromatography by a small computer", *BUNSEKI KAGAKU* **21** (1972) 1043–1051 (CA 78:37709z).
- [63] M. Mitooka, "Gas chromatography with mixed stationary phase. Part XIV. Peak identification with mixed packing gas chromatography", *BUNSEKI KAGAKU* **21** (1972) 1437–1446 (CA 79:26901a).
- [64] M. Mitooka, "Gas chromatography with mixed stationary phase. Part XV. Gas chromatographic separation of saturated hydrocarbons from C₅ to C₈ with a capillary column", *BUNSEKI KAGAKU* **21** (1972) 1447–1456 (CA 79:17588b).
- [65] R. Annino, P.F. McCrea, "Variable selectivity stationary phases for gas chromatography", *ANAL. CHEM.* **42** (1970) 1486–1491.
- [66] V.G. Berezkin, V.E. Smirygaeva, T.P. Popova, "Open-tubular columns with several layers of different immobilized stationary phases", *ZH. ANAL. KHM.* **47** (5) (1992) 825–831 (CA 117:244777j).
- [67] P.P. Nair, I. Sarlos, J. Machiz, "Microquantitative separation of isomeric dimethyltolcols by gas-liquid chromatography", *ARCH. BIOCHEM. BIOPHYS.* **114** (3) (1966) 488–493 (CA 65:2602f).
- [68] H. Bober, "Application of a mixed phase in the gas chromatographic separation of phenols, cresols, and xylenols", *FETTE, SEIFEN, ANSTRICHMITTEL* **68** (6) (1966) 464–468 (CA 65:12885g).
- [69] C. Landault, G. Guiochon, "Fast analysis of phenol, methylphenols, and polymethylphenols by gas chromatography using packed capillary columns", *ANAL. CHEM.* **39** (1967) 713–721.

- [70] D. Solomon, D. Strummer, P.P. Nair, "Routine assay of 11-deoxy-17-keto steroids in human urinary gas-liquid chromatography with a standardized column", *AMER. J. CLIN. PATHOL.* **48** (3) (1967) 295–300 (CA 67:114032a).
- [71] A. Darbre, K. Blau, "Gas chromatography of volatile amino acid derivatives. Part IV. Mixed stationary phases for the separation of n-trifluoroacetylated amino acid n-amyl esters", *J. CHROMATOGR.* **29** (1967) 49–57.
- [72] A. Darbre, A. Islam, "Gas-liquid chromatography of trifluoroacetylated amino acid methyl esters", *BIOCHEM. J.* **106** (4) (1968) 923–925 (CA 68:92766j).
- [73] A. Islam, A. Darbre, "Gas-liquid chromatography of trifluoroacetylated amino acid methyl esters. Development of mixed stationary phase for their separation", *J. CHROMATOGR.* **43** (1969) 11–24.
- [74] A. Andersons, S. Jurels, M.V. Shimanskaya, "Gas-liquid chromatography of some aliphatic and heterocyclic polyfunctional amines. Part III. Separation of mixtures of amines on binary sorbents", *LATV. PSR ZINAT. AKAD. VESTIS, KIM. SER.* (5) (1969) 545–551 (CA 72:96437h).
- [75] A. Pinelli, P.P. Nair, "Gas-liquid chromatographic separation of steroids and their derivatives on a dual component column of high thermal stability", *J. CHROMATOGR.* **43** (1969) 223–228.
- [76] A. Di Muccio, L. Boniforti, R. Monacelli, "Gas chromatographic separation of hexachlorobenzene and the alpha-, beta-, gamma- and delta-isomers of hexachlorocyclohexane", *J. CHROMATOGR.* **71** (1972) 340–346.
- [77] V. Rus, I. Funduc, A. Cralniceanu, S. Trestianu, "Gas chromatographic column for separation of organochlorine insecticide residues", *ANAL. CHEM.* **49** (1977) 2123–2124.
- [78] S.K. Levy, T. Schwartz, "New mixed bed column for rapid gas-liquid chromatographic determination of sedatives and dilantin in serum", *CLIN. CHIM. ACTA* **54** (1) (1974) 19–26 (CA 81:145583z).
- [79] K. van de Castele, H. de Pooter, C.F. van Sumere, "Gas chromatographic separation and analysis of trimethylsilyl derivatives of some naturally occurring nonvolatile phenolic compounds and related substances", *J. CHROMATOGR.* **121** (1976) 49–63 (CA 85:58989a).
- [80] P. Hurter, T. van Ree, "Programmed-temperature gas chromatography on mixed phases. Separation of steroid derivatives on binary mixtures", *J. CHROMATOGR.* **169** (1979) 93–100.
- [81] A. Ono, "Separation of mononitroxylene and xylenol isomers by gas-liquid chromatography", *J. CHROMATOGR.* **180** (1979) 170–176.
- [82] A. Ono, "Separation of xylenol isomers on nitrofluorenes and nitro-9-fluorenones by gas-liquid chromatography. Part I. Mixed columns and effect of temperature on retention behaviour", *J. CHROMATOGR.* **193** (1980) 300–307.
- [83] L. Zoccolillo, G. Cartoni, L. Lozzi, "Binary stationary-phase columns for gas chromatography of barbiturates", *J. CHROMATOGR.* **236** (1982) 339–344.
- [84] J.L. Daft, "Gas chromatographic determination of chemical residues in food using a rugged high-resolution mixed-bed column", *ANAL. CHEM.* **56** (1984) 2687–2692.
- [85] A.C. Butler, R.R. Sibbald, "Investigations into the utility of an inorganic salt eutectic column for trace hydrocarbon determinations", *J. CHROMATOGR. SCI.* **23** (1985) 352–354.
- [86] W. Czelakowski, P. Kusz, A. Andrysiak, "Chromatographic resolution on mixed stationary phases; the choice of the optimum composition", *CHROMATOGRAPHIA* **22** (1986) 278–282.
- [87] G. Castello, T.C. Gerbino, "Optimization of the composition of mixed-phase columns for gas chromatographic separation of C1–C2 halocarbons", *J. CHROMATOGR.* **366** (1986) 59–68.
- [88] A. Bulanova, M.S. Vigdergauz, "Calculation of chromatographic retention of solutes by binary stationary phases based on d-sorbitol", *ZH. ANAL. KHM.* **43** (8) (1988) 1477–1482 (CA 111: 32895x).
- [89] K. Shimakura, H. Yamanaka, K. Shiomi, T. Kikuchi, "Gas liquid chromatographic determination of pyrrolidine and piperidine in canned fish products", *NIPPON SUISAN GAKKAISHI* **57** (9) (1991) 1753–1757 (CA 115:254439y).
- [90] C. Benmei, W. Xiangpu, X. Minru, "Gas chromatographic determination of chlorothalonil concentration in the air of workshops", *ANALYSIS* **20** (5) (1992) 291–293 (CA 117:117349w).
- [91] B. Polanuer, "Direct aqueous injection gas chromatographic analysis of polar compounds using sorbents containing potassium fluoride dihydrate", *CHROMATOGRAPHIA* **33** (1992) 279–283.
- [92] B.M. Polanuer, "Direct aqueous injection gas chromatography on a potassium fluoride crystal hydrate-containing sorbent. Determination of volatile organic solvents in the fermentation broth of clostridium strains", *J. CHROMATOGR.* **596** (1992) 138–140.

- [93] J. Mangas, M.P. Gonzales, D. Blanco, "Influence of cider-making technology on low-boiling-point volatile compounds", *Z. LEBENSM.-UNTERS. FORSCH.* **197** (6) (1993) 522–524 (CA 121: 203852u).
- [94] T.A. Anderson, E.L. Kruger, J.R. Coats, "Enhanced degradation of a mixture of three herbicides in the rhizosphere of a herbicide-tolerant plant", *CHEMOSPHERE* **28** (1994) 1551–1557 (CA 121: 29167n).
- [95] G.M. Kheifets, E.M. Alekseeva, "Gas chromatographic determination of traces of blood plasma cholestryl esters", *ZH. ANAL. KHM.* **49** (3) (1994) 307–314 (CA 120:264978r).
- [96] N. Kaushik, A. Agnihotri, "Evaluation of improved method for determination of rapeseed-mustard fames by GC", *CHROMATOGRAPHIA* **44** (1997) 97–99.
- [97] J.A. Burke, W. Holswade, "Gas chromatographic column for pesticide residue analysis: retention times and response data", *J. ASSOC. OFF. ANAL. CHEM.* **49** (2) (1966) 374–385 (CA 65:2894g).
- [98] J.C. Touchstone, C.H. Wu, A. Nikolski, T. Murawec, "Retention behavior of steroids in gas chromatography with a series of combination columns", *J. CHROMATOGR.* **29** (1967) 235–238.
- [99] F.K. Kawahara, "Microdetermination of derivatives of phenols and mercaptans by means of electron capture gas chromatography", *ANAL. CHEM.* **40** (1968) 1009–1010.
- [100] R.R. Watts, R.W. Storherr, "Gas chromatography of organophosphorus pesticides: retention times and response data on three columns", *J. ASSOC. OFF. ANAL. CHEM.* **52** (3) (1969) 513–521 (CA 71:12019s).
- [101] E.S. Windham, "Gas chromatographic column for pesticide analysis", *J. ASSOC. OFF. ANAL. CHEM.* **52** (6) (1969) 1237–1239 (CA 72:30547n).
- [102] J.F. Thompson, A.C. Walker, R.F. Moseman, "Study of the performance of gas chromatographic columns under severe injection loading", *J. ASSOC. OFF. ANAL. CHEM.* **52** (6) (1969) 1251–1262 (CA 72:30549q).
- [103] J.F. Thompson, A.C. Walker, R.F. Moseman, "Evaluation of eight gas chromatographic columns for chlorinated pesticides", *J. ASSOC. OFF. ANAL. CHEM.* **52** (6) (1969) 1263–1277 (CA 72:30550h).
- [104] J.C. Touchstone, T. Murawec, A. Nikolski, "Retention slopes of steroids on gas chromatography in series of combination columns", *J. CHROMATOGR. SCI.* **8** (1970) 221–223.
- [105] J.C. Touchstone, T. Murawec, "Mixed phase packings of OV substrates for gas chromatography of steroids", *J. CHROMATOGR. SCI.* **9** (1971) 446–447.
- [106] J.A. Armour, "Gas chromatographic data for polychlorinated biphenyl components in six Aroclors", *J. CHROMATOGR.* **72** (1972) 275–282.
- [107] G. Westoo, A. Andersson, "Determination of biphenyl and 2-phenylphenol in citrus fruits by gas-liquid chromatography", *ANALYST* **100** (1975) 173–177 (CA 83:56786s).
- [108] W.A. Prater, M.S. Simmons, K.H. Mancy, "Microanalysis of aqueous samples for phenols and organic acids", *ANAL. LETT.* **13** (A3) (1980) 205–212 (CA 93:79629p).
- [109] T.F. Guerin, S.W.L. Kimber, I.R. Kennedy, "Efficient one-step method for the extraction of cyclodiene pesticides from aqueous media and the analysis of their metabolites", *J. AGRIC. FOOD CHEM.* **40** (11) (1992) 2309–2314 (CA 117:227847q).
- [110] T.F. Guerin, I.R. Kennedy, "Distribution and dissipation of endosulfan and related cyclodienes in sterile aqueous systems: implications for studies on biodegradation", *J. AGRIC. FOOD CHEM.* **40** (11) (1992) 2315–2323 (CA 117:207112t).
- [111] J. Mes, L. Marchand, D. Turton, P.Y. Lau, P.R. Ganz, "The determination of polychlorinated biphenyl congeners and other chlorinated hydrocarbon residues in human blood, serum and plasma", *INT. J. ENVIRON. ANAL. CHEM.* **48** (3-4) (1992) 175–186 (CA 117:246853e).
- [112] G.E. Miliadis, "Gas chromatographic determination of pesticides in natural waters of Greece", *BULL. ENVIRON. CONTAM. TOXICOL.* **50** (2) (1993) 247–252 (CA 118:66373k).
- [113] D. Bourgeois, J. Gaudet, P. Deveau, V.N. Mallet, "Microextraction of organophosphorus pesticides from environmental water and analysis by gas chromatography", *BULL. ENVIRON. CONTAM. TOXICOL.* **50** (3) (1993) 433–440 (CA 118:131628p).
- [114] R.W. Davis, S.T. Kamble, M.P. Tolley, "Microencapsulated chlorpyrifos distribution in loamy sand and silty clay loam soils when applied with a sub-slab injector for subterranean termite control", *BULL. ENVIRON. CONTAM. TOXICOL.* **50** (3) (1993) 458–465 (CA 118:163226x).
- [115] C.J. Kim, L. Taylor, F. Selman, "Effect of herbicide formulation on atrazine residue removal from chemical plant workers' protective clothing", *BULL. ENVIRON. CONTAM. TOXICOL.* **50** (6) (1993) 811–816 (CA 119:14496d).

- [116] D.J. Forsyth, P.A. Martin, "Effects of fenitrothion on survival, behavior, and brain cholinesterase activity of white-throated sparrows (*Zonotrichia Albicollis*)", ENVIRON. TOXICOL. CHEM. **12** (1) (1993) 91–103 (CA 118:75058p).
- [117] K.D. Hoagland, R.W. Drenner, J.D. Smith, D.R. Cross, "Freshwater community responses to mixtures of agricultural pesticides: effect of atrazine and bifenthrin", ENVIRON. TOXICOL. CHEM. **12** (4) (1993) 627–637 (CA 119:33853j).
- [118] B.A. Rattner, M.J. Melancon, T.W. Custer, R.L. Hothem, K.A. King, L.J. LeCaptain, J.W. Spann, B.R. Woodin, J.J. Stegman, "Biomonitoring environmental contamination with piping black-crowned night heron embryos: induction of cytochrome P450", ENVIRON. TOXICOL. CHEM. **12** (9) (1993) 1719–1732 (CA 119:232827d).
- [119] P. Tremolada, D. Calamari, C. Gaggi, E. Bacci, "Fingerprints of some chlorinated hydrocarbons in plant foliage from Africa", CHEMOSPHERE **27** (1993) 2235–2252 (CA 120:85286s).
- [120] C. Albelda, Y. Pico, G. Font, J. Manes, "Determination of aldicarb, aldicarb sulfoxide, and aldicarb sulfone in oranges by simple gas-liquid chromatography with nitrogen-phosphorus detection", J. AOAC INT. **77** (1) (1994) 74–78 (CA 120:189978k).
- [121] G.E. Miliadis, "Determination of pesticide residues in natural waters of Greece by solid phase extraction and gas chromatography", BULL. ENVIRON. CONTAM. TOXICOL. **52** (1) (1994) 25–30.
- [122] P. Aplada-Sarlis, K.S. Liapis, G.E. Miliadis, "Contamination of potato tubers and carrots in Greece with lindane residues", BULL. ENVIRON. CONTAM. TOXICOL. **52** (1) (1994) 135–140.
- [123] M.A. El-Dib, R.K. Ali, "Mixed algal population and *Scenedesmus* Sp. as trihalomethane precursors", BULL. ENVIRON. CONTAM. TOXICOL. **52** (5) (1994) 712–717.
- [124] V. Drevendar, B. Stengl, Z. Frobe, "Microanalysis of dialkyl phosphorus metabolites of organic phosphorus pesticides in human blood by capillary gas chromatography and by phosphorus selective and ion trap detection", ANAL. CHIM. ACTA **290** (1994) 277–286.
- [125] S. Fingler, B. Tkalcovic, Z. Froebe, V. Drevendar, "Analysis of polychlorinated biphenyls, organochlorine pesticides and chlorophenols in rain and snow", ANALYST **119** (1994) 1135–1140 (CA 121:90333e).
- [126] J. Mes, D.L. Arnold, F. Bryce, "Determination of polychlorinated biphenyls in postpartum blood, adipose tissue, and milk from female Rhesus monkeys and their offspring after prolonged dosing with Aroclor 1254", J. ANAL. TOXICOL. **18** (1) (1994) 29–35 (CA 120:127059c).
- [127] R.S. Henly, R.F. Kruppa, W.R. Supina, "Mixed-phase silicone packing for pesticide analysis by gas chromatography", J. AGRIC. FOOD CHEM. **14** (6) (1966) 667–668 (CA 66:10133a).
- [128] C.W. Gehrke, K. Kuo, R.W. Zumwalt, "The complete gas-liquid chromatographic separation of the twenty protein amino acids", J. CHROMATOGR. **57** (1971) 209–217.
- [129] H.H. Nejad, D.W. Kellog, L.B. Daniels, "GC analysis of chlorinated insecticides in bovine feces", CAN. J. ANIM. SCI. **70** (1) (1990) 337–341 (CA 113:1636a).
- [130] A. Tanaka, M. Iijima, Y. Kikuchi, "Determination of hydrogen peroxide in fish products and noodles (Japanese) by gas-liquid chromatography with electron-capture detection", J. AGRIC. FOOD. CHEM. **38** (12) (1990) 2154–2159 (CA 113:229780d).
- [131] S.K. Pant, P.N. Gupta, K.M. Thomas, B.K. Maltin, C.L. Jain, "Simultaneous determination of camphor, menthol, menthyl salicylate, and thymol in analgesic ointments by gas-liquid chromatography", LC-GC INT. **3** (6) (1990) 36–38.
- [132] S.M. Waliszewski, G.A. Szymczynski, "Persistent organochlorine pesticides in blood serum and whole blood", BULL. ENVIRON. CONTAM. TOXICOL. **46** (6) (1991) 803–809 (CA 115:23803w).
- [133] M.L. Ribeiro, E.V. Minelli, "On-line DDT determination in blood serum: experimental parameters", BULL. ENVIRON. CONTAM. TOXICOL. **47** (6) (1991) 804–810 (CA 116:53039q).
- [134] D.K. Verma, "Gas liquid chromatographic analysis of DDT residues in kidney of *Labeo Rohita*", ACTA HYDROCHIM. HYDROBIOL. **19** (3) (1991) 327–329 (CA 115:87148u).
- [135] A. Di Muccio, A. Ausili, R. Dommarco, D. Attard Barbini, A. Santilio, F. Vergori, G. de Merulis, L. Sernicola, "Solid-matrix partition for separation of organochlorine pesticide residues from fatty materials", J. CHROMATOGR. **552** (1991) 241–247 (CA 115:157300p).
- [136] P.L. Eberbach, L.A. Douglas, "Method for the determination of glyphosate and (aminomethyl) phosphonic acid in soil using electron capture gas chromatography", J. AGRIC. FOOD CHEM. **39** (10) (1991) 1776–1780 (CA 115:177302u).
- [137] J.E. France, J.V. King, J.M. Snyder, "Supercritical fluid-based cleanup technique for the separation of organochlorine pesticides from fats", J. AGRIC. FOOD CHEM. **39** (10) (1991) 1871–1874 (CA 115: 181643y).

- [138] B. Hennig, F. Luedcke, "Solid-phase extraction for the determination of isosorbide dinitrate and its metabolites in plasma", GIT. FACHZ. LAB. **35** (11) (1991) 1225–1226 (CA 116:207238t).
- [139] A. Hollstein, "Residues of organochlorine pesticides, PCBs and chlorophenols in fishes from fish-hatcheries, lakes and the river Oder in the area Frankfurt (Oder)", NAHRUNG **35** (10) (1991) 1029–1039 (CA 116:180652j).
- [140] Z. Huang, H. Nie, S. Peng, "Method for determination of residual chlordimeform in food by bromination gas chromatography", SEPU **9** (1) (1991) 53–55 (CA 114:205626c).
- [141] H.M. Lee, C.W. Lee, "Determination of morphine and codeine in blood and bile by gas chromatography with a derivatization procedure", J. ANAL. TOXICOL. **15** (4) (1991) 182–187 (CA 115: 107962k).
- [142] P.P. Singh, R.L. Kalra, "Use of low-purity solvents and reagents for analysis of residues of synthetic pyrethroid pesticides by gas chromatography with electron capture detection", J. CHROMATOGR. **540** (1991) 464–468 (CA 114:159009d).
- [143] C. Yang, Y. Zhang, S. Ren, "Determination of theophylline by gas chromatography ", SEPU **9** (5) (1991) 324–325 (CA 115:189904e).
- [144] G.H. Tan, "Comparison of solvent extraction and solid-phase extraction for the determination of organochlorine pesticide residues in water", ANALYST **117** (1992) 1129–1132 (CA 117:96979j).
- [145] Y. Lu, X. Xue, J. Xu, W. Yan, "Gas chromatographic determination of acaricide danjiami in citrus and soil", J. AOAC INT. **75** (6) (1992) 1100–1102 (CA 118:34346n).
- [146] F. Luedcke, B. Hennig, B. Vetter, "Gas-chromatographic analysis of trace amounts of isosorbide dinitrate and glycerol trinitrate in human plasma after solid-phase extraction", PHARMAZIE **47** (8) (1992) 640–641 (CA 120:22982v).
- [147] J.A. Akkara, M. Greenberger, D.L. Kaplan, "New analytical methods for quantitation of four fungicides by gas and high-performance liquid chromatography", J. CHROMATOGR. **604** (1992) 247–253 (CA 117:123797g).
- [148] Y. Ji, S. Wang, Q. Zhang, W. Li, J. Lu, "Determination of residues of organochlorine, organophosphorous and carbamate pesticides in water by gas chromatography", SEPU **10** (2) (1992) 94–96 (CA 117:55514k).
- [149] J.C. Mugachia, L. Kanja, T.E. Maitho, "Organochlorine pesticide residues in estuarine fish from the Athi river, Kenya", BULL. ENVIRON. CONTAM. TOXICOL. **49** (2) (1992) 199–206 (CA 117: 55450m).
- [150] S.M. Waliszewski, "Residues of lindane, HCH isomers and HCB in the soil after lindane application", ENVIRON. POLLUT. **82** (3) (1993) 289–293 (CA 119:264635f).
- [151] N. Fernandez, M. Sierra, J.J. Garcia, M.J. Diez, M.T. Teran, "Organochlorine pesticide residues in black tea, chamomile, and linden", BULL. ENVIRON. CONTAM. TOXICOL. **50** (4) (1993) 479–485 (CA 118:132109g).
- [152] J. Gomez-Catalan, J. Planas, J. To-Figueras, M. Camps, J. Corbella, "Organochlorine pesticide residues in the population of Catalonia (Spain)", BULL. ENVIRON. CONTAM. TOXICOL. **51** (1) (1993) 160–164 (CA 119:14430c).
- [153] A. Jabbar, S.Z. Masud, Z. Parveen, M. Ali, "Pesticide residues in cropland soils and shallow groundwater in Punjab, Pakistan", BULL. ENVIRON. CONTAM. TOXICOL. **51** (2) (1993) 268–273 (CA 119:79571u).
- [154] G.N. Ramaswamy, C.R. Boyd, "Fluorescent dye and pesticide penetration tested in a computerized spray chamber. Part I. Non-woven fabrics as barriers", BULL. ENVIRON. CONTAM. TOXICOL. **51** (3) (1993) 341–348 (CA 119:145549w).
- [155] A. Arino, A. Herrera, P. Conchello, R. Lazaro, "Hexachlorobenzene and hexachlorocyclohexane residues in pork as affected by weight and sex", BULL. ENVIRON. CONTAM. TOXICOL. **51** (5) (1993) 647–650 (CA 119:224638c).
- [156] A. Katayama, F. Matsumura, "Degradation of organochlorine pesticides, particularly endosulfan, by Trichoderma Harzianum", ENVIRON. TOXICOL. CHEM. **12** (6) (1993) 1059–1065 (CA 119: 67724h).
- [157] D.J. Hoffman, G.J. Smith, B.A. Rattner, "Biomarkers of contaminant exposure in common terns and black-crowned night herons in the Great Lakes", ENVIRON. TOXICOL. CHEM. **12** (6) (1993) 1095–1103 (CA 119:55542j).

- [158] R.W. Chadwick, S.E. George, J. Chang, M.J. Kohan, J.P. Dekker, J.C. Allison, J.E. Long, M.C. Duffy, L.R. Forehand, "Effects of age, species difference, antibiotics and toxicants on intestinal enzyme activity and genotoxicity", *ENVIRON. TOXICOL. CHEM.* **12** (8) (1993) 1339–1352 (CA 120:2353n).
- [159] A. Arino, A. Herrera, M.P. Conchello, R. Lazaro, C. Perez, "Hexachlorocyclohexane residues in meat products after processing", *J. FOOD COMPOS. ANAL.* **6** (1) (1993) 55–61 (CA 119:224642z).
- [160] I. Mukherjee, M. Gopal, "Organochlorine pesticide residues in dairy milk in and around Delhi", *J. AOAC INT.* **76** (2) (1993) 283–286 (CA 118:232566x).
- [161] J.M. Snyder, J.W. King, L.D. Rowe, J.A. Woerner, "Supercritical fluid extraction of poultry tissues containing incurred pesticide residues", *J. AOAC INT.* **76** (4) (1993) 888–892 (CA 120:7048b).
- [162] W.J. Trotter, R. Dickerson, "Pesticide residues in composited milk collected through the U.S. pasteurized milk network", *J. AOAC INT.* **76** (6) (1993) 1220–1225 (CA 120:161995q).
- [163] I. Mukherjee, M. Gopal, "New method for the determination of residues of oxydemeton methyl in mustard crop by gas chromatography of its sulfone", *FRESENIUS' J. ANAL. CHEM.* **347** (1993) 126–128 (CA 120:2694z).
- [164] V. Schurig, J. Zhu, V. Muschalek, "Enantiomer separation on amylose tris(n-butylcarbamate) by gas chromatography", *CHROMATOGRAPHIA* **35** (1993) 237–240.
- [165] R.L. Kalra, B. Singh, R.S. Battu, "Organochlorine pesticide residues in human milk in Punjab, India", *ENVIRON. POLLUT.* **85** (2) (1994) 147–151 (CA 120:296983e).
- [166] H. Steinwandter, "A collaborative study for intermethod comparison. Micro and macro extraction methods for the determination of organophosphorus pesticides in fruits and vegetables", *FRESENIUS' J. ANAL. CHEM.* **348** (1994) 688–691.
- [167] G.H. Tan, K. Vijayaletchumy, "Organochlorine pesticide residue levels in peninsular Malaysian rivers", *BULL. ENVIRON. CONTAM. TOXICOL.* **53** (3) (1994) 351–356.
- [168] G.M. D'Eril, N. Cappuccia, M. Colli, V. Molina, "Gas chromatography of 4,4'-diphenylmethane diisocyanate in workplace atmosphere", *J. CHROMATOGR. A* **718** (1995) 141–146.
- [169] R.E. Leveque, "Determination of C₃–C₈ hydrocarbons in naphthas and reformates utilizing capillary column gas chromatography", *ANAL. CHEM.* **39** (1967) 1811–1818.
- [170] P. Merchant, Jr., "Resolution of C₄ to C₁₂ petroleum mixtures by capillary gas chromatography", *ANAL. CHEM.* **40** (1968) 2153–2158.
- [171] R.G. Mathews, J. Torres, R.D. Schwartz, "Separation of 28 to 114 °C hydrocarbons with glass capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **1** (1978) 139–142.
- [172] P. Sandra, M. van Roelenbosch, "Mixed phases of Superox 20M and OV-1 in fused silica and glass capillary gas chromatography", *CHROMATOGRAPHIA* **14** (1981) 345–350.
- [173] H.T. Badings, C. de Jong, "Headspace analysis for the study of aroma compounds in milk and dairy products", in *ANALYSIS OF VOLATILES. METHODS AND APPLICATIONS* (Proc. Int. Workshop, Würzburg, FRG, September 28–30, 1983), P. Schreier, Ed., Walter de Gruyter, Berlin (1984) 401–417.
- [174] T. Czajkowska, "Glass capillary columns with mixed stationary phases", *J. CHROMATOGR.* **243** (1982) 35–42.
- [175] M. Horka, V. Kahle, M. Krejci, "Immobilized polyethylenglycol-based stationary phase for the analysis of amines and organic acids by capillary gas chromatography", *J. CHROMATOGR.* **637** (1993) 96–99.
- [176] Gy. Vigh, J. Hlavay, Z. Varga-Puchony, T. Welsch, "Preparation and use of high temperature silylated capillary columns coated with a mixture of OV-1 and FFAP for the analysis of volatile carboxylic acids", *J. HIGH RESOLUT. CHROMATOGR.* **5** (1982) 124–127.
- [177] J. Pörschmann, T. Welsch, W. Engewald, Gy. Vigh, "Characterization of OV-1/FFAP-mixture coated glass capillary columns used for the separation of free fatty acids", *J. HIGH RESOLUT. CHROMATOGR.* **7** (1984) 509–514.
- [178] J. Sanz, M. de Frutos, I. Martinez-Castro, "Design and evaluation of a mixed-phase capillary column for the gas chromatographic separation of the volatile components of cheese", *CHROMATOGRAPHIA* **33** (1992) 213–217.
- [179] M. de Frutos, J. Sanz, I. Martinez-Castro, M.I. Jimenez, "New parameters for the characterization of relationship between gas chromatographic retention and temperature", *ANAL. CHEM.* **65** (1993) 2643–2649.

- [180] S. Oi, Y. Ochiai, S. Miyano, "Separation of enantiomeric amino acid derivatives on axially chiral 1,1'-binaphthyl-2,2'-bis(n-decylcarboxamide) as a stationary phase for capillary gas chromatography", CHEM. LETT. (9) (1991) 1575–1576 (CA 115:247167e).
- [181] H.H. Oelert, H.J.K. Koeser, J. Stemmler, "Extended application of gas chromatography for high-boiling and nonvolatile hydrocarbons. Part I. Development, characterization and application of separation columns based on salt melts.", BER. – DTSCH. GES. MINERALWISS. KOHLECHEM. **241** (1981) Pt. 1, 156 pp. (CA 97:8873k).
- [182] M. Taramasso, F. Veniale, CONTR. MINERAL. PETROL. **21** (1959) 1796.
- [183] M. Taramasso, F. Veniale, "Gas chromatographic performance of long-chain alkylammonium complexes with beidellite-type clay minerals", CHROMATOGRAPHIA **2** (1969) 239–242.
- [184] P. Fuchs, M. Taramasso, "Specific interactions and steric effects on beidellite type interlamellar complexes", CHROMATOGRAPHIA **2** (1969) 551–553.
- [185] M. Taramasso, P. Fuchs, "The organic complexes of the clay minerals and their use in gas chromatography", J. CHROMATOGR. **49** (1970) 70–75.
- [186] P. Fuchs, "Organic derivatives of Hungarian bentonites in hydrocarbon analysis", J. CHROMATOGR. **65** (1972) 219–225.
- [187] G.A. Eiceman, A.S. Lara, "Gas chromatographic properties of organoammonium exchanged montmorillonites. Part I. Tetraalkylammonium cations", J. CHROMATOGR. **549** (1991) 273–281.
- [188] A.S. Lara, G.A. Eiceman, "Gas chromatographic properties of organoammonium exchanged montmorillonites. Part II. $\text{CH}_3\text{N}^+\text{H}_3$, $(\text{CH}_3)_4\text{N}^+$, $(\text{CH}_3)_3\text{N}^+\text{CH}_2\text{C}_6\text{H}_5$ and 1,4-diazabicyclo-[2.2.2] octane-2 H^+ ", J. CHROMATOGR. **549** (1991) 283–295.
- [189] A. Zlatkis, J. Jiao, "Evaluation of new organoclay stationary phases for the separation of ethylbenzene and xylene isomers", CHROMATOGRAPHIA **31** (1991) 457–464.
- [190] J.M. Vergnaud, B. Buathier, M. Fatscher, J.C. Zanetton, "Use of Bentone 34 in gas chromatography", J. CHROMATOGR. **27** (1967) 54–62 (CA 66:121860b).
- [191] P. Matousek, "Rapid gas chromatography of isomeric aromatic hydrocarbons", FRESENIUS' Z. ANAL. CHEM. **236** (1) (1968) 201–207 (CA 69:8336j).
- [192] C.F. Raley, J.W. Kaufman, "Separation of chlorostyrenes and ethylbenzene/xylenes", ANAL. CHEM. **40** (1968) 1371–1372.
- [193] H.W. Duerbeck, "Separation of isomeric C₆–C₁₀ aromatic hydrocarbons by gas chromatography", FRESENIUS' Z. ANAL. CHEM. **251** (1970) 108–111 (CA 73:72847m).
- [194] F.W. Karasek, "Analysis of brominated benzene compounds by gas chromatography", J. CHROMATOGR. SCI. **8** (1970) 282–284.
- [195] F.W. Karasek, I. Fong, "Analysis of chlorinated benzene compounds by gas chromatography", J. CHROMATOGR. SCI. **9** (1971) 497–499.
- [196] F.W. Karasek, P. Vassilakopoulos, "Analysis of bromo-chlorobenzenes by gas chromatography", CHROMATOGRAPHIA **4** (1971) 388–390.
- [197] F.W. Karasek, T. Stepanik, "Quantitative analysis of trichlorobenzenes by gas chromatography", J. CHROMATOGR. SCI. **10** (1972) 573–575.
- [198] D. Deur-Siftar, V. Svob, "Separation of xylenes by gas chromatography", KEM. IND. **20** (11) (1971) 549–553 (CA 77:69831d).
- [199] D.W. Grant, "The use of gel-coated glass microspherical support for improved column performance and the rapid separation of isomeric compounds", J. CHROMATOGR. **81** (1973) 33–45.
- [200] D.W. Grant, "Some observation of the use of dimethyldioctadecylammonium bentonite as a stationary phase in gas chromatography", CHROMATOGRAPHIA **6** (1973) 239–240.
- [201] S.-E. Chitour, J.-M. Vergnaud, "A gas-liquid chromatographic study of the adsorption properties of a stationary phase containing a mixture a squalane and Bentone-34", J. CHROMATOGR. **89** (1974) 295–307.
- [202] D.M. Ottenstein, D.A. Bartley, W.R. Supina, "Gas chromatographic separation of styrene in the presence of xylenes and propylbenzenes", ANAL. CHEM. **46** (1974) 2225–2226.
- [203] K. Kawata, Y. Fujieda, "Gas chromatographic determination of volatile hydrocarbons in air by using adsorbent collection", JPN. J. TOXICOL. ENVIRON. HEALTH **38** (4) (1992) 355–360 (CA 117: 197298s).
- [204] E.J. Knust, M. Schueller, "Gas chromatographic separation of mono- and disubstituted benzene derivatives and of diastereoisomeric halogenalkanes with Bentone-34", J. CHROMATOGR. **114** (1975) 207–210.

- [205] Holderith J., Tóth T., Váradi A., „A gázkromatográfiás elemzési idő minimalizálása. Az optimális mérési paraméterek megkeresése szimplex módszer egítségével”, MAGY. KÉM. FOLY. **81** (1975) 162.
- [206] J. Holderith, T. Tóth, A. Váradi, "Minimizing the time for gas chromatographic analysis. Search for optimal operational parameters by a simplex method", J. CHROMATOGR. **119** (1976) 215.
- [207] V.G. Berezkin, V.S. Gavrichev, "Retention of hydrocarbons on gas chromatographic sorbents containing an organo-clay", J. CHROMATOGR. **116** (1976) 9–16.
- [208] F. Vernon, A.N. Khakoo, "Comparison of bentone and liquid-crystal stationary phases for the gas chromatographic separation of isomers", J. CHROMATOGR. **157** (1978) 412–415.
- [209] A. Bhattacharjee, D. Guha, "Gas chromatography of low-boiling pyridine bases", J. CHROMATOGR. **298** (1984) 164–168.
- [210] G.-C. Lin, C.A. Chang, "Heavy alkylated benzene stationary phase for the gas-liquid chromatographic separation of ethylbenzene, xylenes, styrene, iso- and n-propylbenzene", J. CHROMATOGR. **409** (1987) 371–376.
- [211] A.M. Nardillo, R.C. Castells, C.B. Castells, "Analysis of isomeric cresols by gas chromatography", CHROMATOGRAPHIA **32** (1991) 457–460.
- [212] S.A. Arystanbekova, A.I. Zhukova, Y.I. Taresevich, S.V. Bondarenko, Z.M. Rivina, V.I. Rozhkov, "Use of organosubstituted montmorillonite for gas-chromatographic analysis of technical 1,2-dichloroethane", UKR. KHIM. ZH. **57** (7) (1991) 719–722 (CA 116:120099z).
- [213] X. Chen, Z. Xu, Y. Ding, "Determination of trace vinyl chloride, vinylidene chloride and ethylidene dichloride in food packagings by mixed stationary liquid of DNP and organic bentonite. Head space gas chromatography", SEPU **9** (3) (1991) 204–205 (CA 115:181658g).
- [214] H. Kelker, B. Scheurle, H. Winterscheidt, "Flüssig-kristalline schmelzen bei temperaturen unterhalb 100C und oberhalb 200C als stationare phasen in der gaschromatographie", ANAL. CHIM. ACTA **38** (1967) 17-39.
- [215] J.P. Schroeder, D.C. Schroeder, M. Katsikas, "Nematic mixtures as stationary liquid phases in gas-liquid chromatography", in PROC. 2ND AMER. CHEM. SOC. SYMP. LIQUID CRYST. ORDERED FLUIDS (1969), J.F. Johnson, Ed., Plenum, New York (1970) 169–180 (CA 78: 151890b).
- [216] M.A. Andrews, D.C. Schroeder, J.P. Schroeder, "Liquid crystals. Part II. 4,4'-di-n-alkoxyazobenzenes and their mixtures as stationary liquid phases in gas liquid chromatography", J. CHROMATOGR. **71** (1972) 233–242.
- [217] J.E. Haky, G.M. Muschik, "Gas chromatographic properties of mixed liquid crystal stationary phases", J. CHROMATOGR. **238** (1982) 367–375.
- [218] Y. Chen, S. Chen, X. Ma, "Studies on liquid crystals used as stationary phases in gas chromatography. Part II. Chromatographic behavior of some mixed liquid crystals", GAODENG XUEXIAO HUAXUE XUEBAO **4** (2) (1983) 194–200 (CA 99:115259c).
- [219] Y. Chen, "Chromatographic analysis of liquid aromatization products of lower hydrocarbons by using liquid crystal M-103", HUAXUE SHIJIE **26** (12) (1985) 456–458 (CA 104:141480j).
- [220] R.J. Laub, W.L. Roberts, "Use of mixed phases for enhanced gas chromatographic separation of polycyclic aromatic hydrocarbons: preliminary studies with liquid crystals", in 4TH INT. SYMP. POLYNUCL. AROMAT. HYDROCARBONS: CHEM. BIOL. EFF. (1979), A. Bjoerseth, A.J. Dennis, Eds., (1980) 25–28 (CA 95:54346x).
- [221] R.J. Laub, W.L. Roberts, C.A. Smith, "Fabrication of gas-chromatographic glass capillary columns of high efficiency with liquid crystal stationary phases", J. HIGH RESOLUT. CHROMATOGR. **3** (1980) 355–356.
- [222] R.J. Laub, W.L. Roberts, C.A. Smith, "Use of mixed phases for enhanced gas chromatographic separation of polycyclic aromatic hydrocarbons. Part II. Prediction of retentions with open-tubular columns of high efficiency with liquid-crystal solvents", in 5TH INT. SYMP. CHEM. ANAL. AND BIOL. FATE: POLYNUCL. AROMAT. HYDROCARBONS (1980), M. Cooke, A.J. Dennis, Eds., (1981) 287–295 (CA 95:231448e).
- [223] H. Finkelmann, R.J. Laub, W.L. Roberts, C.A. Smith, "Use of mixed phases for enhanced gas chromatographic separation of polycyclic aromatic hydrocarbons. Part III. Phase transition behavior, mass transfer nonequilibrium, and analytical utility of mesogen solvents", REPORT DOE/ER/10554-14-REV.1, Order No. DE82020763 (1982) 15 pp. (CA 98:209346k).

- [224] H. Finkelmann, R.J. Laub, W.L. Roberts, C.A. Smith, "Use of mixed phases for enhanced gas-chromatographic separation of polycyclic hydrocarbons. Part III. Phase transition behavior, mass-transfer nonequilibrium, and analytical properties of a mesogen polymer solvent with silicone diluents", in 6TH INT. SYMP. POLYNUCL. AROMAT. HYDROCARBONS: PHYS. BIOL. CHEM. (1981), M. Cooke, A.J. Dennis, G.L. Fischer, Eds., Battelle Press, Columbus (1982) 275–285 (CA 99:32530h).
- [225] G.M. Janini, G.M. Muschik, H.J. Issaq, R.J. Laub, "Neat and admixed mesomorphic polysiloxane stationary phases for open-tubular column gas chromatography", ANAL. CHEM. **60** (1988) 1119–1124.
- [226] F. Hahne, G. Kraus, H. Zaschke, "Blended liquid crystalline phases in capillary gas chromatography", J. CHROMATOGR. **520** (1990) 85–90.
- [227] R.C. Kong, M.L. Lee, Y. Tominaga, R. Pratap, M. Iwao, R.N. Castle, S.A. Wise, "Capillary column gas chromatographic resolution of isomeric polycyclic sulfur heterocycles in a coal liquid", J. CHROMATOGR. SCI. **20** (1982) 502–510.
- [228] W. Klaue, G. Kraus, "Application of liquid-crystal separation phases for gas chromatographic analysis of organochlorine pesticides and polychlorinated biphenyls", LEBENSMITTEL INDUSTRIE **35** (4) (1988) 156–157 (CA 110:130360t).
- [229] O.K. Guha, J. Janak, "Charge-transfer complexes of metals in the chromatographic separation of organic compounds", J. CHROMATOGR. **68** (1972) 325–343.
- [230] V. Schurig, "Enantiomer analysis by complexation gas chromatography. Scope, merits and limitations", J. CHROMATOGR. **441** (1988) 135–153.
- [231] D. Cagniant, "Complexation chromatography", Chromatographic science series, Vol. 57, D. Cagniant, Ed., Marcel Dekker, New York (1992) (CA 116:222213z).
- [232] R. Villalobos, "Preparation and application of long-lived silver nitrate columns in gas chromatography", ANAL. INSTRUM. 1965, (pub. 1966) 13–19 (CA 71:35763u).
- [233] H.J. Ache, A.P. Wolf, "Radio-gas chromatographic analysis of ¹¹C-labeled isotopic ethylenes and acetylenes", FREZENIUS' Z. ANAL. CHEM. **230** (1) (1967) 19–28 (CA 67:104937k).
- [234] J.G. Atkinson, A.A. Russell, R.S. Stuart, "Gas chromatographic studies of isotopically labeled ethylenes", CAN. J. CHEM. **45** (1967) 1963.
- [235] E. Bendel, B. Fell, W. Gartzen, G. Kruse, "Gas chromatographic determination of unsaturated hydrocarbons by using silver salt packed columns. Part II. Quantitative separation of all bonding isomers of n-octenes, n-hexynes and hexadienes with conjugated and cumulated double bonds", J. CHROMATOGR. **31** (1967) 531–534 (CA 68:74876r).
- [236] T.E. Wessel, J.A. Berson, "Stabilization of a putative cyclohexane-1,4-diyi intermediate elicits an antarafacial cope rearrangement via a stepwise mechanism. Pyrolysis of (R,E)-5-methyl-1,2,6-octatriene to 4-methyl-3-methylene-1,5-heptadiene", J. AM. CHEM. SOC. **116** (1994) 495–505 (CA 120:243803q).
- [237] D.V. Banthorpe, C. Gatford, B.R. Hollebone, "Gas chromatographic separation of olefins and aromatic hydrocarbons using thallium(I) nitrate–glycol as stationary phase", J. GAS CHROMATOGR. **6** (1) (1968) 61–62 (CA 68:84138r).
- [238] A.M. Soto, N. Yanagihara, T. Ogura, "Silver complex of poly(methylphenylsiloxane) as the active component for separation of aliphatic, olefinic, and aromatic hydrocarbons by gas chromatography", ANAL. CHEM. **63** (1991) 1178–1178.
- [239] S.P. Wasik, W. Tsang, "Gas-liquid chromatography separations of hydrocarbons using columns with aqueous solutions of complexing metal ions as stationary phases", ANAL. CHEM. **42** (1970) 1648–1649.
- [240] S.P. Wasik, W. Tsang, "Gas chromatographic determination of partition coefficients of some unsaturated hydrocarbons and their deuterated isomers in aqueous silver nitrate solutions", J. PHYS. CHEM. **74** (1970) 2970–2976 (CA 73:59511c).
- [241] V.G. Berezkin, E.N. Viktorova, V.S. Gavrichov, S.A. Rang, K.R. Kuningas, A.E. Meister, "Gas chromatographic separation of unsaturated polar compounds on inorganic stationary phases containing silver nitrate", J. CHROMATOGR. **552** (1991) 313–318 (CA 115:222404s).
- [242] V.G. Berezkin, E.N. Viktorova, "Use of inorganic salts for gas chromatographic separation of unsaturated compounds", ZH. ANAL. KHIM. **47** (5) (1992) 835–839 (CA 117:263966m).
- [243] A. Zlatkis, I.M.R. de Andrade, "Capillary columns containing silver nitrate substrate for the analysis of olefins", CHROMATOGRAPHIA **2** (1969) 298–303.

- [244] R.L. Pecsok, E.M. Vary, "Gas-solid-liquid chromatography: dispersed metal phthalocyanines as a substrate", *ANAL. CHEM.* **39** (1967) 289–294.
- [245] R.C. Castells, J.A. Catoggio, "Gas chromatographic study of interactions between aliphatic amines and metallic ions", *ANAL. CHEM.* **42** (1970) 1268–1271.
- [246] M.C. Titon, A.M. Nardillo, "Separation of low-boiling pyridine bases by gas chromatography", *J. CHROMATOGR. A.* **699** (1995) 403–407.
- [247] M.C. Titon, F.R. Gonzalez, A.M. Nardillo, "Gas chromatographic separation of low- boiling pyridine bases", *CHROMATOGRAPHIA* **42** (1996) 465–468.
- [248] A. Bhattacharjee, A. Bhaumik, "Analysis of low-boiling isomers of phenols by gas chromatography", *J. CHROMATOGR.* **136** (1977) 328–331 (CA 87:77981z).
- [249] O.K. Guha, A. Bhaumik, "Gas chromatography of the system phenanthrene-anthracene-carbazole on a column of cadmium chloride modified with Carbowax 20M", *J. CHROMATOGR.* **119** (1976) 181–186 (CA 85:71771t).
- [250] V. Schurig, "Resolution of enantiomers and isotopic compositions by selective complexation gas chromatography on metal complexes", *CHROMATOGRAPHIA* **13** (1980) 263–270.
- [251] V. Schurig, W. Buerkle, "Extending the scope of enantiomer resolution by complexation gas chromatography", *J. AM. CHEM. SOC.* **104** (1982) 7573–7580.
- [252] V. Schurig, R. Weber, "Use of glass and fused-silica open tubular columns for the separation of structural, configurational and optical isomers by selective complexation chromatography", *J. CHROMATOGR.* **289** (1984) 321–332.
- [253] V. Schurig, U. Leyrer, R. Weber, "The use of glass and fused silica open tubular columns for the separation of structural, configurational, and optical isomers by selective complexation GC: sec-alcohols and ketones", *J. HIGH RESOLUT. CHROMATOGR.* **8** (1985) 459–464.
- [254] V. Schurig, "Enantiomer analysis by complexation gas chromatography. Scope, merits and limitations", *J. CHROMATOGR.* **441** (1988) 135–153.
- [255] B.T. Golding, P.J. Sellars, A.K. Wong, "Resolution of racemic epoxides on GLC columns containing optically active lanthanoid complexes", *J. CHEM. SOC. CHEM. COMM.* (1977) 570–571.
- [256] W. Wasiak, "New packings for complexation gas chromatography", *INT. CHROMATOGR. LAB.* **6** (SEPT.) (1991) 2–6.
- [257] W. Wasiak, "Chemically bonded chelates as selective complexing sorbents for gas chromatography. Part I. Alkenes", *J. CHROMATOGR.* **547** (1991) 259–268.
- [258] W. Wasiak, "Chemically bonded chelates as selective complexing sorbents for gas chromatography. Part II. Ketones, ethers and nitroalkanes", *J. CHROMATOGR. A.* **653** (1993) 63–69.
- [259] W. Wasiak, "Specific interactions of n-donor adsorbates with n-benzoylthiourea–copper(II) complexes chemically bonded to silica supports", *CHROMATOGRAPHIA* **41** (1995) 107–111.
- [260] E. Smolkova-Keulemansova, "Cyclodextrins as stationary phases in chromatography", *J. CHROMATOGR.* **251** (1982) 17–34.
- [261] S. Li, W.C. Purdy, "Cyclodextrins and their applications in analytical chemistry", *CHEM. REV.* **92** (1992) 1457–1470.
- [262] W.A. Koenig, "Gas chromatographic enantiomer separation with modified cyclodextrins", Hüthig Verlag, Heidelberg (1992).
- [263] V. Schurig, "Enantiomer separation by gas chromatography on chiral stationary phases", *J. CHROMATOGR. A.* **666** (1994) 111–129.
- [264] V. Schurig, H.-P. Nowotny, "Gas-chromatographic separation of enantiomers on stationary phases without metal complexes. Part II. Gas-chromatographic separation of enantiomers on cyclodextrin derivatives", *ANGEW. CHEM.* **102** (9) (1990) 969–986 (ANGEW. CHEM. INT. ED. ENGL. **29** (1990) 939–957) (CA 113:171108x).
- [265] H.-G. Schmarr, A. Mosandl, H.-P. Neukom, K. Grob, "Modified cyclodextrins as stationary phases for capillary GC: consequences of dilution in polysiloxanes", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 207–210.
- [266] M. Jung, D. Schmalzing, V. Schurig, "Theoretical approach to the gas chromatographic separation of enantiomers on dissolved cyclodextrin derivatives", *J. CHROMATOGR.* **552** (1991) 43–57.
- [267] A. Dietrich, B. Maas, G. Brand, V. Karl, A. Kauzinger, A. Mosandl, "Stereoisomeric flavor compounds. Part LX. Diluted modified cyclodextrins as chiral stationary phases. The influence of the polysiloxane solvent", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 769–772.

- [268] I. Hardt, W.A. Koenig, "Diluted versus undiluted cyclodextrin derivatives in capillary gas chromatography and the effect of linear carrier gas velocity, column temperature, and length on enantiomer separation", *J. MICROCOLUMN SEP.* **5** (1993) 35–40.
- [269] C. Bicchi, G. Artuffo, A. D'Amato, V. Manzin, A. Galli, M. Galli, "Cyclodextrin derivatives for the GC separation of racemic mixtures of volatile compounds. Part VI. The influence of the diluting phase on the enantioselectivity of 2,6-di-O-methyl-3-O-pentyl-beta-cyclodextrin", *J. HIGH RESOLUT. CHROMATOGR.* **16** (1993) 209–214.
- [270] C. Bicchi, G. Artuffo, A. D'Amato, V. Manzin, A. Galli, M. Galli, "Cyclodextrin derivatives in GC separation of racemic mixtures of volatiles. Part VI. The influence of diluting on enantioselectivity of 2,6-dimethyl-3-pentyl-beta-cyclodextrin", in *PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 265–276.
- [271] F. Kobor, G. Schomburg, "6-tert-butyldimethylsilyl-2,3-dimethyl-alpha-, beta-, and gamma- cyclodextrins, dissolved in polysiloxanes, as chiral selectors for gas chromatography. Influence of selector concentration and polysiloxane matrix on enantioselectivity", *J. HIGH RESOLUT. CHROMATOGR.* **16** (1993) 693–699.
- [272] A. Dietrich, B. Maas, A. Mosandl, "Diluted modified cyclodextrins as chiral capillary gas chromatographic stationary phases: influence of the polysiloxane solvents on enantioselectivity and column efficiency", *J. MICROCOLUMN SEP.* **6** (1994) 33–42 (CA 121:291622x).
- [273] A. Dietrich, B. Maas, A. Mosandl, "Diluted modified cyclodextrins as chiral stationary phases. Influence of the polysiloxane solvent: heptakis(2,3-di-O-acetyl-6-O-tert-butyldimethylsilyl)-beta-cyclodextrin", *J. HIGH RESOLUT. CHROMATOGR.* **18** (1995) 152–156.
- [274] W.M. Buda, K. Jaques, A. Venema, P. Sandra, "Retention on mixed cyclodextrin/polysiloxane stationary phases in capillary GC. Evidence for the interaction of cyclodextrin and polysiloxane", *FRESENIUS' J. ANAL. CHEM.* **352** (1995) 679–683 (CA 123:328848v).
- [275] W.M. Buda, K. Jaques, A. Venema, P. Sandra, "Retention on mixed cyclodextrin/polysiloxane phases. Evidence for the interaction of cyclodextrin and polysiloxane", in *PROC. 16TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, September 27–30, 1994), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1994) 164–171.
- [276] V. Schurig, H.P. Nowotny, "Separation of enantiomers on diluted permethylated beta-cyclodextrin by high-resolution gas chromatography", *J. CHROMATOGR.* **441** (1988) 155–163, (CA 109:182788n).
- [277] H.P. Nowotny, D. Schmalzing, D. Wistuba, V. Schurig, "Extending the scope of enantiomer separation on diluted methylated β -cyclodextrin derivatives by high-resolution gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **12** (1989) 383–393.
- [278] V. Schurig, M. Jung, D. Schmalzing, M. Schleimer, J. Duvekot, J.C. Buyten, J.A. Peene, P. Mussche, "CGC enantiomer separation on diluted cyclodextrin derivatives coated on fused silica columns", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 470–474.
- [279] Z. Juvanez, G. Alexander, J. Szejtli, *J. HIGH RESOLUT. CHROMATOGR.* **10** (1987) 105.
- [280] W. Keim, A. Koehnes, W. Meltzow, H. Roemer, "Enantiomer separation by gas chromatography on cyclodextrin chiral stationary phases", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1991) 507–529.
- [281] P. Jing, R.-N. Fu, R.-J. Dai, J.-L. Ge, J.-L. Gu, Z. Huang, Y. Chen, "Consequence of diluting modified beta-cyclodextrins in a side-chain crown ether polysiloxane and in a side-chain liquid-crystalline polysiloxane-containing crown ether as stationary phases in capillary gas chromatography", *CHROMATOGRAPHIA* **43** (1996) 546–550.
- [282] A. Di Corcia, D. Fritz, F. Bruner, "Use of graphitized carbon black for linear gas-liquid-solid chromatography of polar low-boiling compounds", *ANAL. CHEM.* **42** (1970) 1500–1504.
- [283] A. Di Corcia, "Gas-liquid-solid chromatography of free acids", *ANAL. CHEM.* **45** (1973) 492–496.
- [284] F. Bruner, P. Ciccioli, G. Crescentini, M.T. Pistolesi, "Role of the liquid phase in gas-liquid-solid chromatography and its influence on column performance. An experimental approach", *ANAL. CHEM.* **45** (1973) 1851–1859.
- [285] A. Di Corcia, "Analysis of phenols by gas-liquid-solid chromatography", *J. CHROMATOGR.* **80** (1973) 69–74.
- [286] F. Bruner, G. Bertoni, G. Crescentini, "Critical evaluation of sampling and gas chromatographic analysis of halogenic hydrocarbons and other organic air pollutants", *J. CHROMATOGR.* **167** (1978) 399–407.
- [287] F. Mangani, F. Bruner, *J. CHROMATOGR.* **289** (1984) 85.

- [288] W.R. Betz, W.R. Supina, *J. CHROMATOGR.* **471** (1989) 105.
- [289] F. Bruner, G. Crescentini, F. Mangani, "Graphitized carbon black: a unique adsorbent for gas chromatography and related techniques", *CHROMATOGRAPHIA* **30** (1990) 565–572.
- [290] L.M. Sidisky, M.V. Robillard, "Carbon layer open tubular capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **16** (1993) 116–119.
- [291] G. Castello, S. Vezzani, P. Moretti, "The selectivity and polarity of carbon layer open tubular capillary columns modified with a polar liquid phase", *J. HIGH RESOLUT. CHROMATOGR.* **17** (1994) 31–36.
- [292] F. Bruner, L. Lattanzi, F. Mangani, M.A. Rezaii, "Graphite layer open tubular (GLOT) columns in gas chromatography and environmental analysis", *CHROMATOGRAPHIA* **38** (1994) 98–108.
- [293] M.A. Rezaii, L. Lattanzi, "Narrow bore silica capillary columns for liquid-modified adsorption gas chromatography", *CHROMATOGRAPHIA* **38** (1994) 235–238.
- [294] J.H. Purnell, "Gas-chromatographic study of chemical equilibria", in *GAS CHROMATOGR., INT. SYMP. ANAL. INSTRUM. DIV. INSTRU. SOC. AMER.* (1966), **6** (1967) 3–20 (CA 70:100149u).
- [295] C.L. Young, "Use of gas chromatography for the determination of thermodynamic properties", *CHROMATOGR. REV.* **10** (2) (1968) 129–158 (CA 73:39370c).
- [296] C.A. Wellington, "Gas chromatographic studies of complexing reactions", *ADVAN. ANAL. CHEM. INSTRUM.* **11** (1973) 237–291 (CA 81:17241z).
- [297] R.J. Laub, R.L. Pecsok, "Study of charge transfer complexation by gas-liquid chromatography", *J. CHROMATOGR.* **113** (1975) 47–68 (CA 83:27031m).
- [298] C.L. de Ligny, "The investigation of complex association by gas chromatography and related chromatographic and electrophoretic methods", *ADV. CHROMATOGR.* **14** (1976) 265–304 (CA 87: 107251f).
- [299] D.G. Gray, "Gas chromatographic measurements of polymer structure and interactions", *PROG. POLYM. SCI.* **5** (1) (1977) 1–60 (CA 87:102675n).
- [300] R.J. Laub, R.L. Pecsok, "Physicochemical application of gas chromatography", John Wiley and Sons, New York (1978) (CA 89:95512t).
- [301] J.E.G. Lipson, J.E. Guillet, "Study of structure and interactions in polymers by inverse gas chromatography", *DEV. POLYM. CHARACT.* **3** (1982) 33–74 (CA 97:128349r).
- [302] C.G. Smith, P.B. Smith, A.J. Pasztor, Jr., M.L. McKelvy, D.M. Meunier, S.W. Froelicher, "Analysis of synthetic polymers and rubbers", *ANAL. CHEM.* **67** (1995) 97R–126R.
- [303] N.A. Katsanos, R. Thede, F. Roubanikantzopoulou, "Diffusion, adsorption and catalytic studies by gas chromatography", *J. CHROMATOGR. A* **795** (1998) 133–184.
- [304] O. Olabisi, "Polymer compatibility by gas-liquid chromatography", *MACROMOLECULES* **8** (1975) 316–322 (CA 83:79849d).
- [305] C.S. Su, D. Patterson, "Determination by gas-liquid chromatography of the polystyrene–poly(vinyl methyl ether) interaction", *MACROMOLECULES* **10** (1977) 708–710 (CA 87:39993x).
- [306] M. Galin, M.C. Rupprecht, "Gas chromatographic study of the interactions in styrene–dimethylsiloxane block copolymers and blends", *MACROMOLECULES* **12** (1979) 506–511 (CA 91: 21286s).
- [307] G. Dipaola-Baranyi, "Thermodynamic miscibility of various solutes with styrene–butyl methacrylate polymers and copolymers", *MACROMOLEKULES* **14** (3) (1981) 683–687 (CA 95:7885n).
- [308] G. Dipaola-Baranyi, P. Degre, "Thermodynamic characterization of polystyrene–poly(butyl methacrylate) blends", *MACROMOLEKULES* **14** (1981) 1456–1460 (CA 95:151282f).
- [309] C.P. Doube, D.J. Walsh, "Studies of poly(vinyl chloride)/solution chlorinated polyethylene blends by inverse gas chromatography", *EUR. POLYM. J.* **17** (1) (1981) 63–69 (CA 94:175707d).
- [310] J. Klein, H. Widdecke, G. Wolter, "Mixed-polymeric stationary phases in gas chromatography", *J. POLYM. SCI., POLYM. SYMP.* **68** (1981) 221–228 (CA 94:72052g).
- [311] J.E.G. Lipson, J.E. Guillet, "Study of structure and interactions in polymers by inverse gas chromatography", *DEV. POLYM. CHARACT.* **3** (1982) 33–74 (CA 97:128349r).
- [312] G. Dipaola-Baranyi, S.J. Fletcher, P. Degre, "Gas chromatographic investigation of poly(vinylidene fluoride)–poly(methyl methacrylate) blends", *MACROMOLECULES* **15** (1982) 885–889 (CA 97: 7110x).
- [313] J.M. Elorza, M.J. Fernandez-Berridi, J.J. Iruin, G.M. Guzman, "The interaction parameters of a PEO/PVA blend by inverse gas chromatography", in *PROC. 28TH IUPAC MACROMOL. SYMP.* (1982), Int. Union Pure Appl. Chem., Oxford (1982) 683 (CA 99:123308h).

- [314] D.J. Walsh, J.S. Higgins, S. Rostami, K. Weeraperuma, "Compatibility of ethylene–vinyl acetate copolymers with chlorinated polyethylenes. Part II. Investigation of the thermodynamic parameters", MACROMOLEKULES **16** (1983) 391–396 (CA 98:108144n).
- [315] C. Zhikuan, D.J. Walsh, "Inverse gas chromatography of one- and two-phase polymer mixtures", EUR. POLYM. J. **19** (6) (1983) 519–524 (CA 99:176639y).
- [316] Z.Y. Al-Saigh, P. Munk, "Study of polymer-polymer interaction coefficients in polymer blends using inverse gas chromatography", MACROMOLECULES **17** (1984) 803–809 (CA 100:157333s).
- [317] J.M. Elorza, M.J. Fdz-Berridi, J.J. Iruin, G.M. Guzman, "Gas chromatographic determination of the interaction parameter of poly(ethylene oxide)/poly(methyl methacrylate) blends", POLYM. ENG. SCI. **24** (4) (1984) 287–290 (CA 100:175690k).
- [318] A.C. Su, J.R. Fried, T. Lorenz, "Study of poly(2,6-dimethyl-1,4-phenylene oxide) blends by inverse gas chromatography", POLYM. MATER. SCI. ENG. **51** (1984) 275–279 (CA 101:172124p).
- [319] A.K. Nandi, B.M. Mandal, S.N. Bhattacharyya, "Miscibility of poly(methyl acrylate) and poly(vinyl acetate): incompatibility in solution and thermodynamic characterization by inverse gas chromatography", MACROMOLEKULES **18** (1985) 1454–1460 (CA 103:37998c).
- [320] G. Dipaola-Baranyi, J. Richer, W.M. Prest, Jr., "Thermodynamic miscibility of polystyrene–poly-(2,6-dimethyl-1,4-phenylene oxide) blends", CAN. J. CHEM. **63** (1) (1985) 223–227 (CA 102: 221531a).
- [321] S. Klotz, R.H. Schuster, H.J. Cantow, "Compatibility of polymers in polymer blends investigated by gas chromatography", MAKROMOL. CHEM. **187** (6) (1986) 1491–1500 (CA 105:79717g).
- [322] B. Riedl, R.E. Prud'homme, "Thermodynamic study of poly(vinyl chloride)–polyester blends by inverse-phase gas chromatography at 120 °C", J. POLYM. SCI., PART B: POLYM. PHYS **24** (11) (1986) 2565–2582 (CA 106:5792b).
- [323] A.C. Su, J.R. Fried, "On the determination of polymer-polymer interaction parameter by inverse gas chromatography", J. POLYM. SCI., PART C: POLYM. LETT. **24** (7) (1986) 343–352 (CA 105: 79722e).
- [324] A.C. Su, J.R. Fried, "Poly(2,6-dimethyl-1,4-phenylene oxide) blends studied by inverse gas chromatography", ADV. CHEM. SER. **211** (MULTICOMP. POLYM. MATER.) (1986) 59–66 (CA 104: 169245k).

Hivatkozások a 2.5. szakaszban

- [1] T.J. Stark, P.A. Larson, R.D. Dandeneau, J. CHROMATOGR. **279** (1983) 31.
- [2] P. Sandra, F. David, M. Proot, G. Diricks, M. Verstappe, M. Verzele, "Selectivity and selectivity tuning in capillary gas chromatography", J. HIGH RESOLUT. CHROMATOGR. **8** (1985) 782–798.
- [3] J.K. Haken, "Polysiloxane stationary phases", J. CHROMATOGR. **73** (1972) 419–448.
- [4] J.K. Haken, "Polysiloxane stationary phases in gas chromatography", J. CHROMATOGR. **141** (1977) 247–288.
- [5] J.K. Haken, "Developments in polysiloxane stationary phases in gas chromatography", J. CHROMATOGR. **300** (1984) 1–77.
- [6] R.A. Keller, "Standard liquid phases. Pandora's box revisited", J. CHROMATOGR. SCI. **11** (1973) 188–190.
- [7] G.W. Pilgrim, R.A. Keller, "Mixed stationary phases", J. CHROMATOGR. SCI. **11** (1973) 206–209.
- [8] J.R. Mann, S.T. Preston, Jr., "Selection of preferred liquid phases", J. CHROMATOGR. SCI. **11** (1973) 216–220.
- [9] S. Hawkes, "Preferred stationary liquids for gas chromatography", J. CHROMATOGR. SCI. **13** (1975) 115–117.
- [10] A. Eskes, F. Dupuis, A. Dijkstra, H. de Clercq, D.L. Massart, "Application of information theory and numerical taxonomy to the selection of gas-liquid chromatography stationary phases", ANAL. CHEM. **47** (1975) 2168–2174.
- [11] H. Rotzsche, "A new type of polar stationary phase with adjustable selectivity coefficient", in GAS CHROMATOGRAPHY, M. van Swaay, Ed., Butterworths, London (1962) 111–121.
- [12] J. Buyten, L. Blomberg, K. Markides, T. Wännman, J. CHROMATOGR. **237** (1982) 465.

- [13] P. Sandra, M. van Roelenbosch, I. Temmerman, M. Verzele, "Characterization, evaluation and in-situ cross-linking of new polar phases in capillary gas chromatography", *CHROMATOGRAPHIA* **16** (1982) 63–68.
- [14] Q. Wu, M. Hetem, C.A. Cramers, J.A. Rijks, "Preparation of thermally stable phenylpolysiloxane fused silica capillary columns. Optimization and evaluation of the deactivation by capillary GC and solid state ^{29}Si NMR", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 811–816.
- [15] P. Petersson, K.E. Markides, D.F. Johnsson, M. Eguchi, S.L. Reese, J. Curtis, B.E. Rossiter, J.S. Bradshaw, M.L. Lee, "A novel strategy for the construction of chiral copolymeric stationary phases for open tubular column chromatography", in *PROC. 13TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 13–16, 1991), P. Sandra, Ed., Hüthig Verlag, Heidelberg (1991) 242–251.
- [16] H. Frank, I. Abe, G. Fabian, "A versatile approach to the reproducible synthesis of functionalized polysilosane stationary phases", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 444–448.
- [17] B.E. Rossiter, S.L. Reese, S. Morgan, A. Malik, J. Bradshaw, M.L. Lee, *J. MICROCOLUMN SEP.* **4** (1992) 521–527.
- [18] W. Blum, R. Aichholz, "High temperature stable CH_3O -terminated poly(diphenyl/dimethyl)- and poly(diphenyl/3,3,3-trifluoropropylmethyl)siloxane copolymer stationary phases for capillary gas chromatography", *J. MICROCOLUMN. SEP.* **5** (1993) 297–302.
- [19] M. Lymann, S.G. Claude, R. Tabacchi, "Synthesis, characterization and evaluation of cyanopropyl siloxane/silarylene copolymers as moderately polar stationary phases for HRGC", in *PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 236–236.
- [20] A. Malik, S.L. Reese, B.A. Johnson, Y. Zhang, B. Haydock, B.E. Rossiter, J. Curtis, J.S. Bradshaw, M.L. Lee, "Cyanophenylsiloxanes. Universal stationary phases for capillary gas chromatography", in *PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 291–298.
- [21] I. Abe, T. Nishiyama, H. Frank, "A method for reproducible preparation of chiral polysiloxanes with regular repeat units as stationary phases for capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **17** (1994) 9–14.
- [22] J.S. Bradshaw, M.L. Lee, A. Malik, G.-L. Yi, W.-B. Li, K.E. Markides, "Rational synthesis of high performance substituted polysiloxane stationary phases for capillary chromatography", in *PROC. 16TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, September 27–30, 1994), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1994) 288–299.
- [23] E.J. Guthrie, J.J. Harland, "Overview of phase development in capillary gas chromatography", *LC-GC INT.* **7** (1994) 130–134.
- [24] M.F. Mehran, W.J. Cooper, R. Lautamo, R.R. Freeman, W. Jennings, "A new bonded stationary phase for the gas chromatographic separation of volatile priority pollutants and chlorinated pesticides", *J. HIGH RESOLUT. CHROMATOGR.* **8** (1985) 715–717.
- [25] T.O. Tiernan, J.H. Garett, J.G. Solch, L.A. Harden, R.M. Lautamo, R.R. Freeman, "New capillary gas chromatography column for the simultaneous isomer-specific determination of 2,3,7,8-TCDD and 2,3,7,8-TCDF", *CHEMOSPHERE* **20** (1990) 1371–1378 (CA 114:114402k).
- [26] T.O. Tiernan, J.H. Garett, J.G. Solch, L.A. Harden, R.M.A. Lautamo, R.R. Freeman, "Use of computerized modeling to develop new capillary GC column stationary phases for optimum separation of chlorinated dioxin/dibenzofuran isomers", in *CAPILLARY CHROMATOGR.*, W.G. Jennings, J.G. Nikelly, Eds., Hüthig Verlag, Heidelberg (1991) 17–38 (CA 115:225544y).
- [27] V. Schurig, D. Schmalzing, U. Muehleck, M. Jung, M. Schleimer, P. Mussche, C. Duvekot, J.C. Buyten, "Gas chromatographic enantiomer separation on polysiloxane-anchored permethyl-beta-cyclodextrin (Chirasil-dex)", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 713–717.
- [28] D. Schmalzing, M. Jung, S. Mayer, J. Rickert, V. Schurig, "Extending the scope of enantiomer separations on Chirasil-Dex by GLC: comparison with permethylated beta-cyclodextrin dissolved in OV-1701", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 723–729.
- [29] M. Jung, V. Schurig, "Enantiomer separation by GC on Chirasil-Dex: systematic study of cyclodextrin concentration, polarity, immobilization, and column stability", *J. MICROCOLUMN. SEP.* **5** (1993) 11–22.

- [30] M Jung, S. Mayer, V. Schurig, "Enantiomer separation by GC, SFC, and CE on immobilized poly-siloxane-bonded cyclodextrins", *LC-GC INT.* **7** (1994) 340–347.
- [31] V. Schurig, "Enantiomer separation by gas chromatography on chiral stationary phases", *J. CHROMATOGR. A.* **666** (1994) 111–129.
- [32] V. Schurig, "Gas chromatographic separation of enantiomers on optically active metal-complex-free stationary phases", *ANGEW. CHEM.* **96** (10) (1984) 733–752 [*ANGEW. CHEM. INT. ED.* **23** (10) (1984) 747–830].
- [33] M.A. Apfel, H. Finkelmann, G.M. Janini, R.J. Laub, B.-H. Luehmann, A. Price, W.L. Roberts, T.J. Shaw, C.A. Smith, "Synthesis and properties of high-temperature mesomorphic polysiloxane solvents: biphenyl- and terphenyl-based nematic systems", *ANAL. CHEM.* **57** (1985) 651–658.
- [34] K.E. Markides, M. Nishioka, B.J. Tarbet, J.S. Bradshaw, M.L. Lee, "Smectic biphenylcarboxylate ester liquid crystalline polysiloxane stationary phase for capillary gas chromatography", *ANAL. CHEM.* **57** (1985) 1296–1299.
- [35] H. Wang, T. Gen, R. Fu, J. Zhu, X. Wei, H. Li, J. Liu, "Gas chromatographic separation of polycyclic aromatic hydrocarbons on a mesomorphic copolysiloxane stationary phase", *J. CHROMATOGR.* **609** (1992) 414–418.
- [36] G. Kraus, J.M. Thierfelder, L. Sojak, "Highly selective liquid crystalline polysiloxane stationary phase for gas chromatographic separation of isomers", *J. CHROMATOGR. A.* **677** (1994) 197–200.
- [37] X.-C. Zhou, C.-Y. Wu, X.-R. Lu, Y.-Y. Chen, "Application of crown ether compounds as gas chromatographic stationary phases", *J. CHROMATOGR. A.* **662** (1994) 203–218.

Hivatkozások a 2.6. szakaszban

- [1] G.W. Pilgrim, R.A. Keller, "Mixed stationary phases", *J. CHROMATOGR. SCI.* **11** (1973) 206–209.
- [2] P. Sandra, F. David, M. Proot, G. Diricks, M. Verstappe, M. Verzele, "Selectivity and selectivity tuning in capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **8** (1985) 782–798.
- [3] P. Sandra, F. David, "Selectivity tuning in capillary gas chromatography", in *MULTIDIMENSIONAL CHROMATOGRAPHY TECHNIQUES AND APPLICATIONS*, Chromatographic Science Series 50, H.J. Cortes, Ed., Marcel Dekker, New York (1990) 145–189.
- [4] J.V. Hinshaw, Jr., L.S. Ettre, "Selectivity tuning of serially coupled open-tubular (capillary) columns in gas chromatography. Part I. fundamental relationships", *CHROMATOGRAPHIA* **21** (1986) 561–572.
- [5] R.E. Kaiser, R.I. Rieder, "Multi-chromatography. Part I.", *LABOR PRAXIS* **9** (1985) 1130–1135 (CA 103:204874t).
- [6] R.E. Kaiser, R.I. Rieder, "Multidimensional chromatography. Part II.", *LABOR PRAXIS* **9** (1985) 1318–1320 (CA 104:141451a).
- [7] R.E. Kaiser, R.I. Rieder, "Multidimensional chromatography. Part III.", *LABOR PRAXIS* **9** (1985) 1465–1472 (CA 104:141119y).
- [8] R.E. Kaiser, R.I. Rieder, "Multidimensional chromatography. Part IV.", *LABOR PRAXIS* **10** (1986) 52–56 (CA 104:218345d).
- [9] R.E. Kaiser, R.I. Rieder, "Multidimensional chromatography. Part V.", *LABOR PRAXIS* **10** (1986) 184–188 (CA 104:179431g).
- [10] R.E. Kaiser, R.I. Rieder, "Multidimensional chromatography. Part VI.", *LABOR PRAXIS* **10** (1986) 328–330 (CA 104:236548b).
- [11] R.E. Kaiser, R.I. Rieder, "Multi-chromatography. Part VII. Problems in quantitative analysis using gas chromatography: changes in specific correction factors according to variations in peak position and shape", *LABOR PRAXIS* **10** (1986) 862–871 (CA 105:164193c).
- [12] R.E. Kaiser, R.I. Rieder, "Multidimensional chromatography. Part VIII.", *LABOR PRAXIS* **10** (1986) 1014–1018 (CA 105:163924e).
- [13] T. Maurer, W. Engewald, "Serial columns in gas chromatography", *LABOR PRAXIS* **14** (1990) 588–594 (CA 113:164692e).
- [14] W. Engewald, T. Knobloch, T. Maurer, "Application of retention indexes in series-coupled gas chromatographic columns. Part I.", *GIT SPEZIAL CHROMATOGRAPHIE* **11** (2) (1991) 102–105 (CA 115:294082h).

- [15] W. Engewald, T. Knobloch, T. Maurer, "Application of retention indexes in series-coupled gas chromatographic columns. Part II.", GIT FACHZ. LAB. 35 (9) (1991) 985–991 (CA 115:294078m).
- [16] R. Villalobos, R. Pearson, "Application of fused silica capillary columns in process gas chromatography", ADV. INSTRUM. **40** (1) (1985) 275–286 (CA 104:81239a).
- [17] R. Villalobos, R. Pearson, "Application of fused silica capillary columns in process gas chromatography", ISA TRANS. **25** (3) (1986) 55–60 (CA 105:99642g).
- [18] F. Clauss, H. Mahler, T. Maurer, "Fifteen years of capillary columns in process gas chromatography (PGC). Part II. Capillary column circuits", LABOR PRAXIS **17** (1993) 38–45 (CA 120:123811u).
- [19] R. Annino, R. Villalobos, "PROCESS GAS CHROMATOGRAPHY. FUNDAMENTALS AND APPLICATIONS", Instrument Society of America, Research Triangle Park, NC, USA (1992).
- [20] R. Villalobos, "Methods development for on-line gas chromatography", J. CHROMATOGR. SCI. **28** (1990) 341–350.
- [21] T. Doran, J.P. Cross, "Single sample analysis of the mixture O₂, N₂, CH₄, CO, CO₂, C₂H₆, C₃H₈, and n-C₄H₁₀", J. GAS CHROMATOGR. **4** (7) (1966) 260–262 (CA 65:11325b).
- [22] G. Castello, "Gas chromatographic determination of alkyllead compounds in aromatic-base fuels", CHIM. IND. **51** (7) (1969) 700–704 (CA 71:126833h).
- [23] M. Ravey, "Mixed bis-lactams. A highly polar stationary phases for the gas chromatography of low molecular weight hydrocarbons", J. CHROMATOGR. SCI. **16** (1978) 79–81.
- [24] G. Castello, T.C. Gerbino, S. Kanitz, "Gas chromatographic analysis of halocarbons in drinking water by headspace extraction and mixed column separation", J. CHROMATOGR. **247** (1982) 263–272.
- [25] G. Castello, T.C. Gerbino, S. Kanitz, "Sensitivity and linearity of headspace and liquid-liquid extraction techniques for gas chromatographic analysis of halocarbons in water", J. CHROMATOGR. **351** (1986) 165–175.
- [26] K. Funazo, M. Tanaka, K. Morita, M. Kamino, T. Shono, H.-L. Wu, "Pentafluorobenzyl p-toluolsulphonate as a new derivatizing reagent for gas chromatographic determination of anions", J. CHROMATOGR. **346** (1985) 215–225.
- [27] Y.I. Korenman, V.N. Fokin, "Identification of volatile phenols in drinking and natural water", GIG. SANIT. (9) (1991) 89–91 (CA 116:27669a).
- [28] T.G. Andronikashvili, L.G. Eprikashvili, N.V. Pirtskhalava, M.G. Adolashvili, "Chromatographic separation of xylene and monochlorotoluene isomers", IZV. AKAD. NAUK GRUZ., SER. KHIM. **17** (2) (1991) 87–92 (CA 116:15106b).
- [29] T.G. Andronikashvili, L.G. Eprikashvili, N.V. Pirtskhalava, G.N. Kirov, V.P. Valtchev, "Gas chromatographic separation of isomeric benzene derivatives using molecular sieves, combined with partition columns", CHROMATOGRAPHIA **38** (1994) 613–616.
- [30] A.A. Chupalov, I.G. Zenkevich, A.P. Molchanov, "GC characteristics of organic reactions. addition of dichlorocarbene to C-C double bonds. precalculation of retention indices for 1,1-dichlorocyclopropanes", in PROC. 16TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, September 27–30, 1994), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1994) 51–60.
- [31] P.L. Gupta, P. Kumar, "Gas chromatographic estimation of individual xylene isomers in straight-run naphtha cuts", ANAL. CHEM. **40** (1968) 992–995.
- [32] M. Taramasso, F. Veniale, "Gas chromatographic performance of long-chain alkylammonium complexes with beidellite-type clay minerals", CHROMATOGRAPHIA **2** (1969) 239–242.
- [33] M. Taramasso, P. Fuchs, "The organic complexes of the clay minerals and their use in gas chromatography", J. CHROMATOGR. **49** (1970) 70–75.
- [34] D. Deur-Siftar, V. Svob, "Separation of xylenes by gas chromatography", KEM. IND. **20** (11) (1971) 549–553 (CA 77:69831d).
- [35] D.G. Gehring, D.J. Barsotti, H.E. Gibbon, "Chlorofluorocarbons alternatives analysis. Part I. The determination of HFC-134a purity by gas chromatography", J. CHROMATOGR. SCI. **30** (1992) 280–284.
- [36] W.F. Wilhite, O.L. Hollis, "The use of porous-polymer beads for analysis of the Martian atmosphere", J. GAS CHROMATOGR. **6** (1968) 84–88.
- [37] S.A. Liebman, D.H. Ahlstrom, T.C. Creighton, G.D. Pruder, R. Averitt, E.J. Levy, "On-line elemental analysis of gas chromatographic effluents", ANAL. CHEM. **44** (1972) 1411–1415.
- [38] S.A. Liebman, D.H. Ahlstrom, C.D. Nauman, R. Averitt, J.L. Walker, E.J. Levy, "On-line elemental-reaction analyzer as an organic analysis system", ANAL. CHEM. **45** (1973) 1360–1364.

- [39] G. Castello, G. D'Amato, "Mixed Porapak columns for the gas chromatographic separation of light hydrocarbons", ANN. CHIM. (Rome) **69** (9–10) (1979) 541–549 (CA 93:88157k).
- [40] L. Huber, "Rapid and simple analysis of natural gas by GC", J. CHROMATOGR. SCI. **21** (1983) 519–520.
- [41] W.V. Ligon, Jr., R.J. May, "Isomer specific analysis of selected chlorodibenzofurans", J. CHROMATOGR. **294** (1984) 87–98.
- [42] W.V. Ligon, Jr., R.J. May, "Determination of selected chlorodibenzofurans and chlorodibenzodioxins using two-dimensional gas chromatography/mass spectrometry", ANAL. CHEM. **58** (1986) 558–561.
- [43] M. Swerev, K. Ballschmiter, "Capillary column coupling for the analysis of environmental patterns of polychlorodibenzodioxins and polychlorodibenzofurans", FREZENIUS' J. ANAL. CHEM. **327** (1987) 50–51.
- [44] M. Swerev, K. Ballschmiter, "Selective separation of 2,3,7,8-tetrachlorodibenzofuran by high-resolution gas chromatography and by alumina column liquid chromatography", ANAL. CHEM. **59** (1987) 2536–2538.
- [45] K. Ballschmitter, R. Bacher, R. Fischer, A. Mennel, U. Riehle, M. Swerev, "The analysis of halogenated biphenyls (HalxB), dibenzodioxins (HalxDD), and dibenzofurans (HalxDF) by HRGC and mass-spectrometric detection (GC/MS)", in PROC. 13TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 13–16, 1991), P. Sandra, Ed., Hüthig Verlag, Heidelberg (1991) 501–525.
- [46] K. Ballschmiter, R. Bacher, A. Mennel, R. Fischer, U. Riehle, M. Swerev, "The determination of chlorinated biphenyls, chlorinated dibenzodioxins, and chlorinated dibenzofurans by GC-MS", J. HIGH RESOLUT. CHROMATOGR. **15** (1992) 260–270.
- [47] L.A. Harden, J.H. Garrett, J.G. Solch, T.O. Tiernan, D.J. Wagel, M.L. Taylor, "Results of comparative evaluations of various fused silica capillary GC columns for retention of tetrachlorinated dibenzo-p-dioxins and dibenzofurans", CHEMOSPHERE **18** (1–6) (1989) 85–91.
- [48] T.O. Tiernan, J.H. Garrett, J.G. Solch, L.A. Harden, R.M. Lautamo, R.R. Freeman, "New capillary gas chromatography column for the simultaneous isomer-specific determination of 2,3,7,8-TCDD and 2,3,7,8-TCDF", CHEMOSPHERE **20** (1990) 1371–1378 (CA 114:114402k).
- [49] S. Bøwadt, B. Larsen, "Improved congener-specific GC analysis of chlorobiphenyls on coupled CP Sil-8 and HT-5 columns", J. HIGH RESOLUT. CHROMATOGR. **15** (1992) 377–380.
- [50] B. Larsen, S. Bøwadt, R. Tilio, S. Facchetti, "Choice of GC column phase for analysis of toxic PCBs", CHEMOSPHERE **25** (1992) 1343–1348 (CA 118:72837f).
- [51] B. Larsen, S. Bøwadt, S. Facchetti, "Separation of toxic congeners from PCB mixtures on two series coupled narrow-bore columns (50m Sil-8 and 25m HT-5)", INT. J. ENVIRON. ANAL. CHEM. **47** (3) (1992) 147–166 (CA 117:105779y).
- [52] B. Larsen, S. Bøwadt, "HRGC separation of PCB congeners. The state-of-the-art", in PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 503–510.
- [53] B.R. Larsen, "HRGC separation of PCB congeners", J. HIGH RESOLUT. CHROMATOGR. **18** (1995) 141–151.
- [54] M.S. Rahman, S. Bøwadt, B. Larsen, "Dual-column GC analysis of Mediterranean fish for 10 organochlorine pesticides and 62 chlorobiphenyls", in PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 815–821.
- [55] M.S. Rahman, S. Bøwadt, B. Larsen, "Dual-column GC analysis of Mediterranean fish for ten organochlorine pesticides and sixty two chlorobiphenyls", J. HIGH RESOLUT. CHROMATOGR. **16** (1993) 731–735.
- [56] S. Bøwadt, B. Johansson, P. Fruekilde, M. Hansen, D. Zilli, B. Larsen, J. de Boer, "Supercritical fluid extraction of polychlorinated biphenyls from lyophilized fish tissue", J. CHROMATOGR. A. **675** (1994) 189–204.
- [57] G.M. Frame, J.W. Cochran, S.S. Bøwadt, "Complete PCB congener distributions for 17 Aroclor mixtures determined by 3 HRGC systems optimized for comprehensive, quantitative, congener-specific analysis", J. HIGH RESOLUT. CHROMATOGR. **19** (1996) 657–668.
- [58] R.C. Kong, M.L. Lee, Y. Tominaga, R. Pratap, M. Iwao, R.N. Castle, "Mesogenic polysiloxane stationary phase for high-resolution gas chromatography of isomeric polycyclic aromatic compounds", ANAL. CHEM. **54** (1982) 1802–1806.

- [59] G. Castello, A. Timossi, T.C. Gerbino, "Analysis of haloalkanes on wide-bore capillary columns of different polarity connected in series", *J. CHROMATOGR.* **522** (1990) 329–343.
- [60] K.K. Gaines, W.H. Chatham, S.O. Farwell, "Comparison of the SCD and FPD for HRGC determination of atmospheric sulfur gases", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 489–493.
- [61] P. Clair, M. Tua, H. Simian, "Capillary column in series for GC analysis of volatile organic pollutants in atmospheric and alveolar air", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 383–387.
- [62] J. Hamill, T.G. Kee, "The detection of aerosol propellants in body fluids and tissue by gas chromatography–mass spectrometry", *J. FORENSIC SCI. SOC.* **31** (3) (1991) 301–307 (CA 116:1844d).
- [63] K. Yamaguchi, S. Nishi, "GC/MS determination of dichloroethylene isomers in water ", *BUNSEKI KAGAKU* **40** (6) (1991) T131–T136 (CA 115:56790t).
- [64] D. Tanzer, K.G. Heumann, "Gas chromatographic trace-level determination of volatile organic sulfides and selenides and of methyl iodide in Atlantic surface water", *INT. J. ENVIRON. ANAL. CHEM.* **48** (1) (1992) 17–31 (CA 117:76125z).
- [65] J.Y.K. Lai, E. Matisova, D. He, E. Singer, H. Niki, "Evaluation of capillary gas chromatography for the measurement of C2–C10 hydrocarbons in urban air samples for air pollution research", *J. CHROMATOGR.* **643** (1993) 77–90.
- [66] K. Abrahamsson, A. Ekdahl, "Gas chromatographic determination of halogenated organic compounds in water and sediment in the Skagerrak", *J. CHROMATOGR.* **643** (1993) 239–248.
- [67] L. Lattanzi, M.A. Rezaii, "A very efficient column for the separation of 60 volatile organochlorine priority pollutants and for other environmental problems", *CHROMATOGRAPHIA* **38** (1994) 114–117.
- [68] L. Lattanzi, M. Attaran Rezaii, L. Sampaolo, "Evaluation of GC graphite-coated capillary columns from polarity parameters", *CHROMATOGRAPHIA* **40** (1995) 729–733.
- [69] C. Giachetti, G. Zanolo, G.R. Verga, F. Perovanni, A. Assandri, "Gas chromatographic determination of some glycol ethers and glycol ether acetates by FID, O-FID, and MS detection. Preliminary application in simultaneous monitoring of these chemicals in biological matrices", *J. HIGH RESOLUT. CHROMATOGR.* **19** (1996) 383–391.
- [70] W. Keim, A. Koehnes, W. Meltzow, H. Roemer, "Enantiomer separation by gas chromatography on cyclodextrin chiral stationary phases", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1991) 507–529.
- [71] V. Karl, H.-G. Schmarr, A. Mosandl, "Simultaneous stereoanalysis of 2-alkyl-branched acids, esters and alcohols using a selectivity-adjusted column system in multi-dimensional gas chromatography", *J. CHROMATOGR.* **587** (1991) 347–350.
- [72] J.P. Kutter, T.J. Class, "Diastereoselective and enantioselective chromatography of the pyrethroid insecticides allethrin and cypermethrin", *CHROMATOGRAPHIA* **33** (1992) 103–112.
- [73] M. Oehme, R. Kallenborn, K. Wiberg, C. Rappe, "Simultaneous enantioselective separation of chlordanes, a nonachlor compound, and o,p'-DDT in environmental samples using tandem capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **17** (1994) 583–588.
- [74] B. Shi, R.A. Keogh, B.H. Davis, "Gas chromatographic separation of deuterated and optical isomers of di-2-butyl ethers", *J. CHROMATOGR. A.* **678** (1994) 97–102.
- [75] G.P. Cartoni, G. Goretti, B. Monticelli, M.V. Russo, "Evaluation of capillary gas chromatographic columns in series. Analytical application to lemon oil", *J. CHROMATOGR.* **370** (1986) 93–101.
- [76] G.P. Cartoni, G. Goretti, M.V. Russo, "Capillary columns in series for the gas chromatographic analysis of essential oils", *CHROMATOGRAPHIA* **23** (1987) 790–795.
- [77] N. Chanegriha, A. Baaliouamer, "Evaluation of series-coupled gas chromatographic capillaries of different polarities. Application to the resolution of problem pairs of constituents in Algerian cypress essential oil", *J. CHROMATOGR.* **633** (1993) 163–168.
- [78] N. Chanegriha, A. Baaliouamer, C. Rolando, "Polarity changes during capillary gas chromatographic and gas chromatographic mass spectrometric analysis using serially coupled columns of different natures and temperature programming. Application to the identification of constituents of essential oils", *J. CHROMATOGR. A.* **819** (1998) 61–65.
- [79] K.G. Miller, C.F. Poole, T.M.P. Chichila, "Solvent-assisted supercritical fluid extraction for the isolation of semivolatile flavor compounds from the cinnamons of commerce and their separation by series-coupled column gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **18** (1995) 461–471.

- [80] A. Jayatilaka, S.K. Poole, C.F. Poole, T.M.P. Chichila, "Simultaneous micro steam distillation/solvent extraction for the isolation of semivolatile flavour compounds from cinnamon and their separation by series coupled-column gas chromatography", *ANAL. CHIM. ACTA* **302** (1995) 147–162.
- [81] A Marti, P. Georges, "Gas chromatographic determination of volatile components in spirits using coupled columns", *MITT. GEB. LEBENSMITTELUNTERS. HYG.* **83** (5) (1992) 549–553 (CA 119:47749s).
- [82] A. Antonelli, M. Motta, "Determination of volatiles in spirits using a combination of stationary phases in capillary GC", in *PROC. 16TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, September 27–30, 1994), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1994) 692–692.
- [83] A. Antonelli, M. Galli, "Determination of volatiles in spirits using combined stationary phases in capillary GC", *CHROMATOGRAPHIA* **41** (1995) 722–725.
- [84] R.E. Pauls, G.J. Weight, P.S. Munowitz, "A comparison of methods to determine benzene in gasoline boiling range material", *J. CHROMATOGR. SCI.* **30** (1992) 32–39.
- [85] J. Feldman, M. Orchin, "Determination of methyl tert. butyl ether (MTBE) in gasoline", *ANAL. LETT.* **26** (2) (1993) 357–365 (CA 118:127902f).
- [86] W.A. Rubey, R.C. Striebich, M.D. Tissandier, D.A. Tirey, S.D. Anderson, "Gas chromatographic measurement of trace oxygen and other dissolved gases in thermally stressed jet fuel", *J. CHROMATOGR. SCI.* **33** (1995) 433–437.
- [87] C.L. Wilkins, G.N. Giss, R.L. White, G.M. Brissey, E.C. Onyiriuka, "Mixture analysis by gas chromatography/Fourier transform infrared spectrometry/mass spectrometry", *ANAL. CHEM.* **54** (1982) 2260–2264.
- [88] W.H. Hwang, R.J. Clark, W.T. Cooper, "Gas chromatographic separation of PF₃ substituted group 6 metal carbonyl complexes using serially-coupled capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **10** (1987) 504–509.
- [89] L. Castellani, G. Goretti, B. Neri, M.V. Russo, P. Zacchei, "Coupled in series microcapillary gas chromatographic columns betel extract analysis", *ANNALI DI CHIMICA* (Rome) **82** (3–4) (1992) 137–148 (CA 116:251357e).
- [90] J.H. Purnell, P.S. Williams, "General theory for correction for compressibility effects in binary coupled gas chromatographic columns and the procedure for window diagram optimization of relative lengths", *J. CHROMATOGR.* **321** (1985) 249–254.
- [91] J.H. Purnell, M. Rodriguez, P.S. Williams, "Approximate approach to correction for carrier compressibility effects in coupled packed gas chromatographic columns", *J. CHROMATOGR.* **323** (1985) 402–405.
- [92] J.H. Purnell, M. Rodriguez, P.S. Williams, "Experimental verification of the theory of serially coupled gas chromatographic columns", *J. CHROMATOGR.* **358** (1986) 39–51.
- [93] J.H. Purnell, J.R. Jones, M.H. Wattan, "Theory and practice of coupled gas chromatographic columns", *J. CHROMATOGR.* **399** (1987) 99–109.
- [94] J.H. Purnell, M.H. Wattan, "Separation of aliphatic hydrocarbon mixtures by gas chromatography using serial liquid-phase and solid-phase columns", *ANAL. CHEM.* **63** (1991) 1261–1264.
- [95] J.H. Purnell, M.H. Wattan, "Theoretical and experimental comparison of serially linked and mixed-packing gas-liquid chromatography columns", *J. CHROMATOGR.* **555** (1991) 173–182.
- [96] J.H. Purnell, P.S. Williams, G.A. Zabierek, "Optimization of solvent choice and analysis in G.L.C.", in *PROC. 4TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Bad Duerkheim, FRG, 1981), R.E. Kaiser, Ed., Institute of Chromatography, Bad Duerkheim (1981) 573–599 (CA 98:27057j).
- [97] J.H. Purnell, P.S. Williams, "Compressibility effects in the optimization of serially connected gas chromatographic capillary columns by the window diagram technique", *J. HIGH RESOLUT. CHROMATOGR.* **6** (1983) 569.
- [98] J.H. Purnell, P.S. Williams, "Relative retention in serially connected binary gas chromatographic capillary column systems and the implications for window diagram optimization of such systems", *J. CHROMATOGR.* **292** (1984) 197–206.
- [99] J.R. Jones, J.H. Purnell, "Prediction of retention times in serially linked open-tubular gas chromatographic columns and optimization of column lengths", *ANAL. CHEM.* **62** (1990) 2300–2306.
- [100] J.H. Purnell, P.S. Williams, "Theory of the efficiency of serially connected gas chromatographic columns and the effect on optimized speed of analysis", *J. CHROMATOGR.* **325** (1985) 1–12.

- [101] J. Krupcik, G. Guiochon, J.M. Schmitter, "Measurement of retention data on open tubular columns coupled in series", *J. CHROMATOGR.* **213** (1981) 189–201.
- [102] J. Krupcik, J. Mocak, A. Simova, J. Garaj, G. Guiochon, "Optimization of experimental conditions for the analysis of complex mixtures by gas chromatography", *J. CHROMATOGR.* **238** (1982) 1–12.
- [103] J. Krupcik, D. Repka, T. Hevesi, J. Mocak, G. Kraus, "Optimization of stationary phase selectivity for the gas-liquid chromatographic separation of C8 cyclic and aromatic hydrocarbons using squalane and liquid crystal glass capillary columns in series", *J. CHROMATOGR.* **355** (1986) 99–105.
- [104] E. Matisova, E. Kavacicova, J. Garaj, G. Kraus, "Optimization of the separation of multicomponent mixtures by HRGC", *CHROMATOGRAPHIA* **27** (1989) 494–498.
- [105] R. Villalobos, R. Annino, "Computer-aided selection and optimization of chromatographic columns and conditions for multicolumn analysis procedures", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 764–773.
- [106] R. Villalobos, R. Annino, "Computer aided selection and optimization of chromatographic columns and conditions for multicolumn analysis procedures", in *PROC. 11TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Monterey, CA, USA, May 14–17, 1990) P. Sandra, G. Redant, M. Leuridan, Eds., Hüthig Verlag, Heidelberg (1990) 436–465.
- [107] R. Villalobos, R. Annino, "Computer-aided design and optimization of an on-line gas chromatographic procedure for the analysis of purgeable compounds in waste water", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 681–685.
- [108] R. Annino, R. Villalobos, "Rapid computer aided solutions to complex GC application problems", in *PROC. 18TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 20–24, 1996), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1996) 279–294.
- [109] R. Villalobos, "A window diagram for key component analysis in on-line gas chromatography", in *PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 115–122.
- [110] R. Villalobos, "A window diagram for key component analysis in on-line gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **18** (1995) 343–347.
- [111] D.F. Ingraham, C.F. Schoenmaker, W. Jennings, "Optimization of liquid phase mixtures", *J. CHROMATOGR.* **239** (1982) 39–50.
- [112] G. Takeoka, H.M. Richard, M. Mehran, W. Jennings, "Optimization via liquid phase mixtures in capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **6** (1983) 145–151.
- [113] M.F. Mehran, W.J. Cooper, W. Jennings, "The use of coupled columns for the capillary gas chromatographic separation of water pollutants", *J. HIGH RESOLUT. CHROMATOGR.* **7** (1984) 215–217.
- [114] M.F. Mehran, W.J. Cooper, R. Lautamo, R.R. Freeman, W. Jennings, "A new bonded stationary phase for the gas chromatographic separation of volatile priority pollutants and chlorinated pesticides", *J. HIGH RESOLUT. CHROMATOGR.* **8** (1985) 715–717.
- [115] K. Yamaoka, T. Nakagawa, T. Mori, T. Uno, "Additivity of moment vectors for the coupled column in a linear gas chromatography", *BUNSEKI KAGAKU* **24** (2) (1975) 84–88 (CA 83:103683t).
- [116] R.G. Williams, H.D. Mitchell, "Computer spreadsheet calculation of the optimum temperature and column lengths for serially coupled capillary gas chromatographic columns", *J. CHROMATOGR.* **541** (1991) 59–74.
- [117] T.C. Gerbino, G. Castello, "Separation of complex mixtures of industrial solvents by using capillary columns connected in series", in *PROC. 13TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 13–16, 1991), P. Sandra, Ed., Hüthig Verlag, Heidelberg (1991) 889–901.
- [118] T.C. Gerbino, G. Castello, "Separation of mixtures of solvents with wide bore capillary columns connected in series", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 428–433.
- [119] T.C. Gerbino, G. Castello, "Theoretical calculation of the retention of chlorobenzenes on non-polar and polar capillary columns connected in series", *J. CHROMATOGR.* **626** (1992) 207–214.
- [120] X. Lou, X. Liu, L. Zhou, "Series coupled capillary columns for the separations of N,(O)-trifluoroacetyl isopropyl derivatives of D,L-aspartic acid and L-hydroxyproline by gas chromatography", *J. CHROMATOGR.* **634** (1993) 281–288.
- [121] R.E. Murphy, R.R. Heath, J.G. Dorsey, "The optimization of capacity and efficiency when coupling fused silica open tubular columns in gas chromatography", *CHROMATOGRAPHIA* **37** (1993) 65–72.

- [122] D.R. Deans, I. Scott, "Gas chromatographic columns with adjustable separation characteristics", ANAL. CHEM. **45** (1973) 1137–1141.
- [123] D.R. Deans, "A new technique for heart cutting in gas chromatography", CHROMATOGRAPHIA **1** (1968) 18–22.
- [124] V.A. Ezrets, M.S. Vigdergauz, "Control of the selectivity of a composite chromatographic column", ZAVOD. LAB. **43** (3) (1977) 277–281 (CA 86:177858n).
- [125] V. Pretorius, T.W. Smuts, J. Moncrieff, "Control of relative retention in gas chromatography by temperature variation if series-coupled columns", J. HIGH RESOLUT. CHROMATOGR. **1** (1978) 200–201.
- [126] R.E. Kaiser, R.I. Rieder, "Change of polarity by temperature variation of series-coupled micropacked GC columns, measured in retention index data", J. HIGH RESOLUT. CHROMATOGR. **1** (1978) 201–202.
- [127] P.R. Boshoff, T.W. Smuts, "Control of relative retention in gas chromatography by temperature adjustment of series coupled columns: evaluation and application to the analysis of pesticides", J. CHROMATOGR. SCI. **18** (1980) 315–323.
- [128] P.R. Boshoff, "Control of relative retention in gas chromatography by temperature adjustment of series-coupled columns: the role of linear temperature programming", J. CHROMATOGR. SCI. **19** (1981) 238–244.
- [129] R.E. Kaiser, R.I. Rieder, "Polarity change in capillary GC by serial-column temperature optimization (SECAT-mode in cap. GC)", in PROC. 3RD INTERNATIONAL SYMPOSIUM ON GLASS CAPILLARY CHROMATOGRAPHY (Hindelang, Germany, April 29–MAY 3, 1979) R.E. Kaiser, Ed., Institute of Chromatography, Bad Duerkheim (1979) 189–208.
- [130] R.E. Kaiser, R.I. Rieder, "Polarity change in capillary GC by serial-column temperature optimization (SECAT mode in capillary GC)", J. HIGH RESOLUT. CHROMATOGR. **2** (1979) 416–422.
- [131] R.E. Kaiser, "Separation cassettes in gas chromatography. A new technique for both easy and complex analysis, with particular reference to environmental analysis", CHROMATOGRAPHIA **7** (1974) 688–695.
- [132] R.E. Kaiser, L. Leming, L. Blomberg, R.I. Rieder, "Dramatic selectivity changes in series coupled capillaries of differing polarities by independently influencing the residence time in the two capillaries", J. HIGH RESOLUT. CHROMATOGR. **8** (1985) 92–94.
- [133] R.E. Kaiser, R.I. Rieder, L. Leming, L. Blomberg, P. Kusz, "Dramatic selectivity changes on carrier gas flow changes in a series coupled GC capillary tandem", in PROC. 6TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 14–16, 1985) P. Sandra, W. Bertsch, Eds., Hüthig Verlag, Heidelberg (1985) 638–647.
- [134] R.E. Kaiser, R.I. Rieder, L. Leming, L. Blomberg, P. Kusz, "Dramatic selectivity changes on carrier gas flow changes in a series-coupled GC capillary tandem", J. HIGH RESOLUT. CHROMATOGR. **8** (1985) 580–584.
- [135] T. Tóth, H. van Cruchten, J. Rijks, "Tuning of column polarity and peak position in complex chromatograms by means of coupled columns", in PROC. 6TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 14–16, 1985) P. Sandra, W. Bertsch, Eds., Hüthig Verlag, Heidelberg (1985) 769–771.
- [136] T. Tóth, J.A. Rijks, J.H.M. van Cruchten, "A novel application of coupled tandem determination GC columns for the tuning of column polarity and for the determination of RI values in multicomponent mixtures. A new approach in naphtha analysis" in PROC. 5TH DANUBE SYMPOSIUM ON CHROMATOGRAPHY (Yalta, USSR, November 11–16, 1985), Nauka (1985) 169–169.
- [137] J.V. Hinshaw, Jr., L.S. Ettre, "Selectivity tuning of serially coupled open-tubular (capillary) columns in gas chromatography. Part I. Fundamental relationships", CHROMATOGRAPHIA **21** (1986) 561–572.
- [138] J.V. Hinshaw, Jr., L.S. Ettre, "Selectivity tuning of serially connected open-tubular (capillary) columns in gas chromatography Part II. Implementation", CHROMATOGRAPHIA **21** (1986) 669–680.
- [139] L.S. Ettre, J.V. Hinshaw, Jr., "Multi chromatography. Part I. Flow tuning of serially connected capillary GC columns", in PROC. 37TH PITTSBURGH CONF. ANAL. CHEM. APPL. SPECTROSCOPY (Atlantic City, NJ, USA, March 10–14, 1986) Paper No. 1100.
- [140] R.E. Kaiser, R.I. Rieder, "Diethylene glycol in wine", J. HIGH RESOLUT. CHROMATOGR. **8** (1985) 863–866.

- [141] R.E. Kaiser, R.I. Rieder, "Native ethylene glycol in wine: application of a dead volume free very fast deans heart-cut-system on-line multi chromatography", in PROC. 8TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 19–21, 1987) P. Sandra, Ed., Hüthig Verlag, Heidelberg (1987) 531–539.
- [142] R.E. Kaiser, R.I. Rieder, "Native ethylene glycol in wine: application of dead volume free, very fast deans heart-cut system on-line with multi-chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **10** (1987) 240–243.
- [143] C.J. Barinaga, S.O. Farwell, "Noncryogenic FSOT column separation of sulfur-containing gases", *J. HIGH RESOLUT. CHROMATOGR.* **10** (1987) 538–543 (CA 108:48403j).
- [144] W. Engewald, T. Maurer, A. Schiefke, "Investigation of isomeric hydrocarbons by gas-solid chromatography on graphitized thermal carbon black", *PURE APPL. CHEM.* **61** (1989) 2001–2004.
- [145] W. Engewald, R. Reinhardt, A. Steinborn, J. Rohloff, R. Herzschuh, "Gas chromatographic behaviour of isomeric tricyclo[10.4.0.0^{4,9}]hexadecanes", *CHROMATOGRAPHIA* **34** (1992) 253–258.
- [146] A. Steinborn, W. Engewald, R. Reinhardt, "The simultaneous use of different separation modes in multidimensional capillary gas chromatography", in PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 892–900.
- [147] L. Lin, G. Li, "Study of chromatographic behavior with single furnace double carrier gas inlets and an in-series-coupled capillary column system", *SEPU* **7** (4) (1989) 196–200 (CA 112:111178q).
- [148] L. Lin, "A refitted model 102 G-D serially coupled capillary gas chromatography system with a single oven and double carrier gas", *SEPU* **9** (1) (1991) 5–10 (CA 114:156126d).
- [149] J. Efer, T. Maurer, W. Engewald, "Application of selectivity tuning in series-coupled capillary GC columns for the exact quantification of the quenching effect in FPD detection", *CHROMATOGRAPHIA* **29** (1990) 115–119.
- [150] W.M. Buda, K. Jaques, A. Venema, P. Sandra, "Retention on mixed cyclodextrin/polysiloxane phases. Evidence for the interaction of cyclodextrin and polysiloxane", in PROC. 16TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, September 27–30, 1994), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1994) 164–171.
- [151] W.M. Buda, K. Jaques, A. Venema, P. Sandra, "Retention on mixed cyclodextrin/polysiloxane stationary phases in capillary GC. Evidence for the interaction of cyclodextrin and polysiloxane", *FRESENIUS' J. ANAL. CHEM.* **352** (1995) 679–683 (CA 123:328848v).
- [152] M. Jung, D. Schmalzing, V. Schurig, "Theoretical approach to the gas chromatographic separation of enantiomers on dissolved cyclodextrin derivatives", *J. CHROMATOGR.* **552** (1991) 43–57.
- [153] E. Benicka, I. Spanik, D. Takacova, M. Grena, J. Krupcik, P. Sandra, "Study of retention behavior of selected amino acid enantiomers by HRGC with modified cyclodextrins", in PROC. 18TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 20–24, 1996), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1996) 76–83.
- [154] J. Krupcik, M. Grena, I. Spanik, E. Benicka, J. Hrouzek, I. Skacani, P. Sandra, "Computerized optimization of selectivity for direct capillary gas chromatographic multicomponent separations of enantiomers", *J. CHROMATOGR. A.* **779** (1997) 253–262.
- [155] A.V. Bulanova, O.N. Bezrukova, "Retention in gas chromatography when there is a large pressure drop in different sections of a coupled column", *RUSS. CHEM. BULL.* **45** (1996) 2792–2799.
- [156] T.W. Smuts, K. de Clerk, T.G. du Toit, T.S. Buys, "Retention time and effective separation factor in series-coupled columns with different stationary phases", *J. HIGH RESOLUT. CHROMATOGR.* **3** (1980) 124–128.
- [157] T.S. Buys, T.W. Smuts, "The effect of the temperature and pressure on the retention time in series-coupled gas chromatographic columns", *J. HIGH RESOLUT. CHROMATOGR.* **3** (1980) 461–470.
- [158] T.S. Buys, T.W. Smuts, "A study of the effect of temperature and pressure on the retention time in series-coupled columns under conditions of constant mass flow", *J. HIGH RESOLUT. CHROMATOGR.* **4** (1981) 102–108.
- [159] T.S. Buys, T.W. Smuts, "The effect of the temperature dependence of mobile phase viscosity on the retention time in series-coupled gas chromatographic columns", *J. HIGH RESOLUT. CHROMATOGR.* **4** (1981) 317–322.
- [160] T.S. Buys, J.B. Wagener, "Retention time in series-coupled columns: An experimental investigation of non-thermodynamic effects", *J. HIGH RESOLUT. CHROMATOGR.* **5** (1982) 662–667.

- [161] T. Tóth, F. Garay, "Polarity tuning of a satellite capillary tandem system by temperature using two independent GC equipment", in PROC. 8TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 19–21, 1987) P. Sandra, Ed., Hüthig Verlag, Heidelberg (1987) 585–590.
- [162] T. Tóth, F. Garay, "Application of the SATELLITE GC capillary tandem column system for analysis of organic solvents of industrial hygienic interest", in PROC. 6TH DANUBE SYMPOSIUM ON CHROMATOGRAPHY (Varna, Bulgaria, October 12–17. 1987), Jusautor, p. 161.
- [163] F. Garay, T. Tóth, "Creation and application of multi-chromatographic data base by temperature tuned tandem capillary GC column system", 8TH INTERNATIONAL SYMPOSIUM: ADVANCES AND APPLICATION OF CHROMATOGRAPHY IN INDUSTRY (Bratislava, Czechoslovakia, July 2–7. 1989), Abstracts Volume II., C43.
- [164] T. Tóth, "Theory and application of tandem capillary column system of tunable selectivity", 8TH INTERNATIONAL SYMPOSIUM: ADVANCES AND APPLICATION OF CHROMATOGRAPHY IN INDUSTRY (Bratislava, Czechoslovakia, July 2–7. 1989), Abstracts Volume I., Mo L8 (oral lecture).
- [165] T. Tóth, F. Garay, Cs. Lovász, "A simple coupling device for flow tuning of serially coupled flexible fused silica capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 840–842.
- [166] F. Garay, T. Tóth, "Application of serially coupled fused silica capillary column system tuned by flow for organic solvent analysis of industrial hygienic interest", in PROC. 13TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 13–16, 1991), P. Sandra, Ed., Hüthig Verlag, Heidelberg (1991) 818–832.
- [167] T. Tóth, F. Garay, "A novel technique for solving the problem of peak matching in CGC analysis of complex mixtures on two stationary phases of different polarity", in PROC. 9TH DANUBE SYMPOSIUM ON CHROMATOGRAPHY (Budapest, Hungary, August 23–27, 1993).
- [168] Gy.L. Török, J. Endrődi, F. Garay, "Novel peak identification method in gas chromatography using flow tunable serially coupled capillary column system", BALATON SYMPOSIUM '97 ON HIGH-PERFORMANCE SEPARATION METHODS, (Siófok, Hungary, September 3–5, 1997), Book of Abstracts, P-145.
- [169] Garay F., Tóth T., „Multi-kromatográfiás adatbázis létrehozása és alkalmazása hőmérséklet hangolt tandem kapilláris kolonna rendszerrel”, XIV. KROMATOGRÁFIÁS VÁNDORGYŰLÉS ÉS TOVÁBBKÉPZŐ TANFOLYAM (Balatonszáplak, 1989. szeptember 4–7).
- [170] Tóth T., Garay F., Lovász Cs., „Egyszerű nyomáshangolható multi-gázkromatográfiás rendszer megvalósítása és gyakorlati alkalmazása I.”, XV. KROMATOGRÁFIÁS VÁNDORGYŰLÉS ÉS TOVÁBBKÉPZŐ TANFOLYAM (Bükfürdő, 1990. október 8–11.).
- [171] Garay F., Tóth T., Lovász Cs., „Egyszerű nyomáshangolható multi-gázkromatográfiás rendszer megvalósítása és gyakorlati alkalmazása II.”, XV. KROMATOGRÁFIÁS VÁNDORGYŰLÉS ÉS TOVÁBBKÉPZŐ TANFOLYAM (Bükfürdő, 1990. október 8–11.).
- [172] Garay F., Tóth T., „Oldószer légszennyezők gázkromatográfiás azonosítása multi-gázkromatográfiás adatbázissal”, ANALITIKAI NAPOK '91 (Budapest, 1991. január 22–23.) (orális előadás).
- [173] Garay F., Tóth T., Gerlei B., „Gázkromatográfiás modell nyomáshangolható tandem kapilláris oszloprendszerre és újszerű alkalmazása sokkomponenses bonyolult elegyek elemzésére”, XVI. KROMATOGRÁFIÁS VÁNDORGYŰLÉS ÉS TOVÁBBKÉPZŐ TANFOLYAM (Balatonaliga, 1991. november 11–13.).
- [174] Török Gy.L., Endrődi J., Garay F., „Újszerű gázkromatográfiás komponensazonosítási technika áramlással hangolható sorasan kapcsolt kapilláris oszloprendszer segítségével”, XIX. KROMATOGRÁFIÁS VÁNDORGYŰLÉS (Budapest, 1997. június 18–19.).
- [175] D. Repka, J. Krupcik, E. Benicka, J. Garaj, T. Maurer, W. Engewald, "Optimization of selectivity by tuning column temperatures and/or intermediate column pressure for series-coupled columns in a dual-oven GC system", in PROC. 10TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, May 22–25, 1989), P. Sandra, G. Redant, Eds., Hüthig Verlag, Heidelberg (1989) 682–696.
- [176] D. Repka, J. Krupcik, E. Benicka, P.A. Leclercq, J.A. Rijks, "Optimization of selectivity by tuning column temperatures for series-coupled capillary columns in dual-oven gas chromatographic systems", *J. CHROMATOGR.* **463** (1989) 243–251.
- [177] D. Repka, J. Krupcik, E. Benicka, T. Maurer, W. Engewald, "Optimization of selectivity by tuning column temperatures for series-coupled columns in a dual-oven GC system", *J. HIGH RESOLUT. CHROMATOGR.* **13** (1990) 333–337.

- [178] T. Hevesi, J. Krupcik, P. Sandra, "Use of chromatographic models for computerized optimization of coupling-point pressure in dual-column gas chromatography", *J. CHROMATOGR.* **517** (1990) 161–173.
- [179] E. Benicka, J. Krupcik, P. Kuljovsky, D. Repka, J. Garaj, "Computer-assisted optimization of selectivity by tuning temperature and carrier gas pressure drop in two GC capillary columns coupled in series", *MIKROCHIM. ACTA* (1990) III. 1–10.
- [180] E. Benicka, J. Krupcik, D. Repka, P. Kuljovsky, "Threshold criteria used for the optimization of selectivity by tuning intermediate pressure for series-coupled columns in a dual-oven system", *ANAL. CHEM.* **62** (1990) 985–990.
- [181] E. Benicka, J. Krupcik, D. Repka, P. Sandra, "Optimization of selectivity by tuning column temperature of identical capillary columns in series in a dual-oven GC system", *CHROMATOGRAPHIA* **33** (1992) 463–466.
- [182] T. Hevesi, J. Krupcik, J.R. Chretien, "Optimization of separation of multicomponent samples analysed by gas chromatography with three open-tubular columns coupled in series", in *PROC. 15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 155–164.
- [183] T. Hevesi, J. Krupcik, J.R. Chretien, "Modeling the gas chromatographic separation of multi-component samples on a system of three open-tubular columns coupled in series", *FRESENIUS' J. ANAL. CHEM.* **352** (1995) 643–648.
- [184] T. Maurer, TH. Welsch, W. Engewald, "Non-linearity of the plot of log (adjusted retention time) versus carbon number for n-alkanes in series-coupled gas chromatographic columns", *J. CHROMATOGR.* **471** (1989) 245–249.
- [185] W. Engewald, T. Maurer, "Enhanced possibilities for identification by the use of series-coupled capillary gas chromatographic columns. Part I. General exposition and application of the retention index concept", *J. CHROMATOGR.* **520** (1990) 3–13.
- [186] T. Maurer, W. Engewald, A. Steinborn, "Enhanced possibilities for identification using series-coupled capillary gas chromatographic columns. Part II. Retention indices as an identification tool in selectivity tuning", *J. CHROMATOGR.* **517** (1990) 77–86.
- [187] W. Engewald, T. Knobloch, G. Haufe, M. Mueller, V. Pohris, "A novel method for terpene pattern determination of essential oils by selectivity tuning in GC", *FRESENIUS' J. ANAL. CHEM.* **341** (1991) 641–643.
- [188] H.T. Mayfield, S.N. Chesler, "Optimization of selectivity using sequentially coupled capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **8** (1985) 595–601.
- [189] H.T. Mayfield, S.N. Chesler, "Optimization of selectivity using sequentially coupled capillary columns", in *PROC. 6TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 14–16, 1985) P. Sandra, W. Bertsch, Eds., Hüthig Verlag, Heidelberg (1985) 772–785.
- [190] Z. Jianying, "Theory for the selection of optimum temperature and carrier gas velocity in two series-coupled capillary GC system", *J. HIGH RESOLUT. CHROMATOGR.* **10** (1987) 418–420.
- [191] G. Guiochon, J.E.N. Gutierrez, "Theoretical prediction of plate height in serially coupled capillary columns", *J. CHROMATOGR.* **406** (1987) 3–10.
- [192] L.H. Wright, J.F. Walling, "Prediction of retention times in temperature-programmed multichromatography", *J. CHROMATOGR.* **540** (1991) 311–322.
- [193] T.C. Gerbino, G. Castello, "Prediction of retention values in temperature programming of capillary columns of different polarity connected in series", *15TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 24–27, 1993), P. Sandra, G. Devos, Eds., Hüthig Verlag, Heidelberg (1993) 60–67.
- [194] T.C. Gerbino, G. Castello, "Prediction of programmed temperature retention times on linked capillary columns of different polarity", *J. HIGH RESOLUT. CHROMATOGR.* **17** (1994) 597–602.
- [195] J.M. Fougnion, C. David, A. Crocq, C. Genty, "Pentaparametric optimization of resolution in gas chromatography. Case of two serially connected columns", *J. CHROMATOGR.* **556** (1991) 287–304.
- [196] M. Akard, R. Sacks, "Column bifurcation with tunable selectivity for high-speed gas chromatography", *J. CHROMATOGR. SCI.* **31** (1993) 297–304.
- [197] M. Akard, R. Sacks, "Pressure-tunable selectivity for high-speed gas chromatography", *ANAL. CHEM.* **66** (1994) 3036–3041.

- [198] M. Akard, R.D. Sacks, "High-speed GC air monitor using cryointegration for sample collection", *J. CHROMATOGR. SCI.* **32** (1994) 499–505.
- [199] R. Sacks, M. Akard, "High-speed GC analysis of VOCs: tunable selectivity and column selection. Part II.", *ENVIRON. SCI. TECHNOL.* **28** (9) (1994) 428A–433A.
- [200] M. Akard, R. Sacks, "Separation widow diagrams for high-speed GC", *ANAL. CHEM.* **67** (1995) 2733–2737.
- [201] M. Akard, R. Sacks, "Simultaneous temperature and holdup time ratio optimization for tandem column tunable selectivity with high-speed GC", *ANAL. CHEM.* **68** (1996) 1474–1479.
- [202] E.B. Bakeas, P.A. Siskos, "A four factor simplex optimization of serially coupled open tubular columns gas chromatographic system", *J. HIGH RESOLUT. CHROMATOGR.* **19** (1996) 277–283.
- [203] J. Krupcik, D. Repka, T. Hevesi, D. Benicka, J. Garaj, "On the use of sequential simplex procedure for optimization of basic parameters influencing temperature in LTPGC analysis of multicomponent samples", *CHROMATOGRAPHIA* **27** (1989) 367–370.
- [204] E.B. Bakeas, P.A. Siskos, "Effect of temperature programming and pressure on separation number and height equivalent to a theoretical plate in optimization of a serially coupled, open tubular columns gas chromatographic system", *ANAL. CHEM.* **68** (1996) 4468–4473.
- [205] H. Smith, R. Sacks, "Vector model of multiphase separation", *ANAL. CHEM.* **69** (1997) 145–151.
- [206] H. Smith, R. Sacks, "Pressure-tunable GC columns with electronic pressure control", *ANAL. CHEM.* **69** (1997) 5159–5164.
- [207] W. Chinghai, H. Frank, W. Guanghua, Z. Liangmo, E. Bayer, L. Peichang, "Determination of amino acid enantiomers by two-column gas chromatography with valveless column switching", *J. CHROMATOGR.* **262** (1963) 352–359.
- [208] G. Schomburg, H. Husmann, E. Huebinger, W.A. Koenig, "Multidimensional capillary gas chromatography. Enantiomer separation of selected cuts using a chiral second column", *J. HIGH RESOLUT. CHROMATOGR.* **7** (1984) 404–410.
- [209] W. Engewald, R. Reinhardt, A. Steinborn, "The use of shape selective stationary phases in GC. Part I. Cyclodextrin phases", *PURE APPL. CHEM.* **66** (1993) 2253–2256.
- [210] P. Hennig, A. Steinborn, W. Engewald, "Investigation of the composition of Pinus Peuce needle oil by GC-MS and GC-GC-MS", *CHROMATOGRAPHIA* **38** (1994) 689–693.
- [211] A. Steinborn, R. Reinhardt, W. Engewald, K. Wyssuwa, K. Schulze, "Enantiomer separation of alpha-campholene and fencholene derivatives by capillary gas chromatography on permethylated cyclodextrins. Part II. Compounds separable with coupled techniques", *J. CHROMATOGR. A.* **697** (1995) 485–494.
- [212] J.A. Rijks, J.H.M. van den Berg, J.P. Diependaal, "Characterization of hydrocarbons in complex mixtures by two-dimensional precision gas chromatography", *J. CHROMATOGR.* **91** (1974) 603–612.
- [213] F. Etzweiler, N. Neuner-Jehle, "A simple effluent stream-splitting device for high resolution glass capillary columns", *CHROMATOGRAPHIA* **6** (1973) 503.
- [214] E.L. Anderson, W. Bertsch, "Practical aspects of Pt/Ir effluent splitters for multidetector GC and pneumatic solute switching", *J. HIGH RESOLUT. CHROMATOGR.* **1** (1978) 13–17.
- [215] F. Rinderknecht, B. Wenger, "The use of platinum-iridium for connecting or interfacing capillary columns in gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **2** (1979) 746–748.
- [216] K. Grob, *CHROMATOGRAPHIA* **9** (1976) 509.
- [217] H.T. Slover, "Use of gold-plated Pt-Ir tubing for joining glass capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **6** (1983) 392–394.
- [218] D. Mueller, H. Walther, "All-glass capillary T-joints and manifolds", *J. HIGH RESOLUT. CHROMATOGR.* **3** (1980) 411–412.
- [219] J.R. Hudson, S.L. Morgan, "Glass capillary T-connections for high resolution GC", *J. HIGH RESOLUT. CHROMATOGR.* **4** (1981) 186–187.
- [220] V. Pretorius, P. Apps, W. Bertsch, "Micro-glass working for capillary chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **6** (1983) 104–105.
- [221] F. Etzweiler, "Flexible capillary effluent splitter of the all-glass type", *J. HIGH RESOLUT. CHROMATOGR.* **7** (1984) 578–579.
- [222] A.-B. Wassgren, G. Bergstroem, "A simple effluent splitter for capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **7** (1984) 154–155.
- [223] V. Pretorius, "A useful cement for gas chromatographers", *J. HIGH RESOLUT. CHROMATOGR.* **3** (1980) 23–24.

- [224] D.W. Later, B.W. Wright, M.L. Lee, "Construction of an efficient fused silica capillary column effluent splitter for gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **4** (1981) 406–408.
- [225] P. Sandra, M. Schelfaut, M. Verzele, "A simple method for connecting fused silica and glass capillaries", *J. HIGH RESOLUT. CHROMATOGR.* **5** (1982) 50–51.
- [226] V. Pretorius, P.J. Apps, E.R. Rohwer, K.H. Lawson, "A means of improving the performance of heat-shrinkable, polytetrafluoroethylene joints in capillary gas liquid chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **8** (1985) 77–78.
- [227] G. Alexander, B.R. Gandhe, "Improved heat-shrinkable PTFE joints also for thin capillaries", *J. HIGH RESOLUT. CHROMATOGR.* **10** (1987) 156–157.
- [228] A. Spark, "A simple, universal joining system for glass to silica capillary tubing", *J. HIGH RESOLUT. CHROMATOGR.* **9** (1986) 481–482.
- [229] R.I. Rieder, R.E. Kaiser, "A simple high temperature, high pressure safe T-piece for installing micro packed glass columns as well as macrobore capillaries in a GC instrument", *J. HIGH RESOLUT. CHROMATOGR.* **9** (1986) 595–595.
- [230] E.R. Rohwer, V. Pretorius, "Hybrid fused silica/glass capillary columns for GLC", *J. HIGH RESOLUT. CHROMATOGR.* **10** (1987) 145–147.
- [231] M. Vecchi, W. Walther, "Simple and versatile method for connecting fused silica and glass capillaries", *J. HIGH RESOLUT. CHROMATOGR.* **11** (1988) 337–338.
- [232] B. Schilling, K. Grob, P. Pichler, R. Dubs, B. Brechbuehler, "Fused-silica ends for glass capillaries: widening glass capillary butts by a heated tungsten point for preparing press-fit connections", *J. CHROMATOGR.* **435** (1988) 204–209.
- [233] K. Grob, Jr., "Connecting capillaries by fusion", *J. CHROMATOGR.* **330** (1985) 217–226.
- [234] P. Sandra, M. Verzele, E. Vanluchene, "Construction of a polyimide effluent splitter for multiple detection in capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **6** (1983) 504–507.
- [235] J. Roeraade, S. Blomberg, G. Flodberg, "Miniature connector with a low dead volume for fused-silica capillary columns", *J. CHROMATOGR.* **301** (1984) 454–460.
- [236] E.R. Rohwer, V. Pretorius, P.J. Apps, "Simple press-fit connector for flexible fused silica tubing in gas-liquid chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **9** (1986) 295–297.
- [237] A. Bemgard, C. Oestman, "High temperature and high pressure stable gluing of press-fit connectors for fused silica and metal capillary tubing", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 131–133.
- [238] C. Wesén, H. Mu, "Re-use of press-fit connectors and splitters for GC capillary columns", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 136–136.
- [239] B. Xu, S. Hu, "A simple technique for making all-glass manifolds for capillary gas chromatography", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 775–778.
- [240] C. Ibanez, "Construction of an all-glass/fused silica, zero dead volume multidimensional gas chromatographic system using press-fit connectors: preliminary results", *J. HIGH RESOLUT. CHROMATOGR.* **16** (1993) 552–554.
- [241] „Multidimensional chromatography. SiCHROMAT 2 the GC with two ovens”, SIEMENS TECHNICAL BROCHURE, Siemens AG, Measurement and Control Division.
- [242] F. Mueller, M. Oreans, "New horizons in high performance gas chromatography using a live-controlled column-switching system", SIEMENS ANALYTICAL APPLICATION NOTE No. 282.
- [243] F. Mueller, M. Oreans, F. Weeke, "Novel column switching according to the live principle", LABOR PRAXIS 6 (1982) 462–472 (CA 97:103633r).
- [244] G. Schomburg, F. Weeke, F. Mueller, M. Oreans, "Multidimensional gas chromatography (MDC) in capillary columns using double oven instruments and a newly designed coupling piece for monitoring detection after pre-separation", *CHROMATOGRAPHIA* **16** (1982) 87–91.
- [245] M. Oreans, F. Mueller, D. Leonhardt, "Trace-analysis in coupled systems; total transfer of traces from packed to capillary columns", in PROC. 5TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Riva del Garda, Italy, April 26–28, 1983), J. Rijks, Ed., Elsevier, Amsterdam (1983) 461–473.
- [246] M. Oreans, F. Mueller, D. Leonhardt, "Trace analysis in coupled systems. Total transfer of traces from packed to capillary columns", *J. CHROMATOGR.* **279** (1983) 357–364.
- [247] F. Mueller, "Column switching in capillary gas chromatography", *AM. LAB.* **15** (10) (1983) 94–103 (CA 99:182058z).

- [248] U. Goekeler, "Multidimensional, double-oven gas chromatography for jet fuel analysis", AM. LAB. **16** (10) (1984) 66–73 (CA 101:232821a).
- [249] F. Mueller, U.K. Goekeler, "Multidimensional, high resolution GC using a double oven", AM. LAB. **17** (5) (1985) 25–38 (CA 103:31789p).
- [250] B.M. Gordon, C.E. Rix, M.F. Borgerding, "Comparison of state-of-the-art column switching techniques in high resolution gas chromatography", J. CHROMATOGR. SCI. **23** (1985) 1–10.
- [251] „MUSIC – The separation of complex mixtures”, CHROMPACK TECHNICAL BROCHURE, Chromapack Int. B.V.
- [252] „SGE chromatography products” SGE catalogue 1992/93, SGE Co., p. 132.
- [253] „Analysis of chiral components in flavours using the MDS-2000”, SGE APPLICATION NOTE, SGE Co. (1993).
- [254] „MCS – Multi column switching gas chromatographic system”, GERSTEL TECHNICAL BROCHURE, Gerstel GmbH.
- [255] „SGE chromatography products” SGE catalogue 1992/93, SGE Co., p. 134.
- [256] F. Mueller, M. Oreans, "Experience and problems with capillary glass columns in process chromatography", CHROMATOGRAPHIA **10** (1977) 473–477.
- [257] W. Dosch, "Unit construction system for coupling operations in high resolution gas chromatography", in PROC. 4TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY (Bad Duerkheim, FRG, 1981), R.E. Kaiser, Ed., Institute of Chromatography, Bad Duerkheim (1981) 549–561.
- [258] H. Broetell, G. Rietz, S. Sandqvist, M. Berg, H. Ehrsson, "Two-dimensional capillary gas chromatography without intermediate trapping. Electron capture detector quantitation of an amino alcohol (Kabi 2128) in serum after trifluoroacetylation", J. HIGH RESOLUT. CHROMATOGR. **5** (1982) 596–603.
- [259] S. Nitz, F. Drawert, U. Gellert, "A variable effluent splitter for simultaneous sniffing-ms-monitoring", CHROMATOGRAPHIA **22** (1986) 51–54.
- [260] Lovász Cs., Garay F., „Egy újszerű csatolóelem multidimenziós kapilláris gázkromatográfiás oszloprendszerek megvalósításához”, VEGYÉSZKONFERENCIA (Eger, 1996. július 2–4.), Előadásösszefoglalók, poszterek.
- [261] Endrődi J., Török Gy.L., Lovász Cs., Garay F., „Újszerű kapilláris gázkromatográfiás csatolóelem alkalmazása multidimenziós és GC-MS rendszerekben”, XIX. KROMATOGRÁFIÁS VÁNDORGYŰLÉS (Budapest, 1997. június 18–19.).
- [262] J. Endrődi, Gy.L. Török, Cs. Lovász, F. Garay, "Applying a novel coupling device to multidimensional capillary gas chromatography and GC-MS", BALATON SYMPOSIUM'97 ON HIGH-PERFORMANCE SEPARATION METHODS, (Siófok, Hungary, September 3–5, 1997), Book of Abstracts, P-94.
- [263] Garay F., Lovász Cs., „MDSS Millennium: újszerű Deans-típusú kapilláris gázkromatográfiás csatlakoztatóelem multidimenziós és GC-MS rendszerekhez”, XX. KROMATOGRÁFIÁS VÁNDORGYŰLÉS (Budapest, 1999. július 7–8.).
- [264] Cs. Lovász, F. Garay, "MDSS Millennium: A novel Deans-type coupling element for multidimensional capillary gas chromatography", BALATON SYMPOSIUM'99 ON HIGH-PERFORMANCE SEPARATION METHODS, (Siófok, Hungary, September 1–3, 1999), Book of Abstracts, P-49.

Hivatkozások a 3. szakaszban

- [1] R.L. Gatrell, "A mixed-substrate column for gas chromatographic analysis of lacquer thinners", ANAL. CHEM. **35** (1963) 923–924.
- [2] K. Kuwata, K. Yamada, H. Yamasaki, Y. Yamazaki, Y. Kiso, "Rapid determination of solvent components in paint by computer-aided fused-silica capillary gas chromatography", J. CHROMATOGR. **256** (1983) 303–312.
- [3] J. Saito, I. Masayuki, "Solvent constituents in paint, glue, and thinner for plastic miniature hobby", TOHOKU J. EXP. MED. **155** (3) (1988) 275–283 (CA 110:9747w).
- [4] Y. Higashino, "Analysis of components of paint solvent by multidimensional gas chromatography", SHIKIZAI KYOKAISHI **62** (12) (1989) 737–743 (CA 113:61347s).

- [5] I.D. Smith, D.G. Waters, "Determination of residual solvent level in bulk pharmaceuticals by capillary gas chromatography", *ANALYST* **116** (1991) 1327–1331.
- [6] T.C. Gerbino, G. Castello, "Separation of complex mixtures of industrial solvents by using capillary columns connected in series", in *PROC. 13TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 13–16, 1991), P. Sandra, Ed., Hüthig Verlag, Heidelberg (1991) 889–901.
- [7] S.C. Rastogi, "Analysis of organic solvents in printing inks by headspace gas chromatography–mass spectrometry", in *PROC. 13TH INTERNATIONAL SYMPOSIUM ON CAPILLARY CHROMATOGRAPHY* (Riva del Garda, Italy, May 13–16, 1991), P. Sandra, Ed., Hüthig Verlag, Heidelberg (1991) 1387–1394.
- [8] S.C. Rastogi, "Analysis of organic solvents in printing inks by headspace gas chromatography–mass spectrometry", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 587–589.
- [9] G. Castello, S. Vezzani, T.C. Gerbino, "Gas chromatographic separation and automatic identification of complex mixtures of organic solvents in industrial wastes", *J. CHROMATOGR.* **585** (1991) 273–280.
- [10] Z. Penton, "Determination of residual solvent in pharmaceutical preparations by static headspace GC", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 329–331.
- [11] T.C. Gerbino, G. Castello, "Separation of mixtures of solvents with wide bore capillary columns connected in series", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 428–433.
- [12] S.C. Rastogi, "Gas chromatographic analysis of organic solvent mixtures on capillary columns of different polarity", *CHROMATOGRAPHIA* **37** (1993) 211–214.
- [13] C. Woolley, J. Murnyack, C. Kepler, "Organic volatile impurities in pharmaceutical products: selectivity of capillary GC columns", *SUPELCO INT.* **14** (2) (1995) 2–3.
- [14] "Resolving industrial solvent mixtures on packed and capillary GC columns", *SUPELCO GC BULLETIN* 824A.
- [15] J.Q. Walker, D.L. Ahlberg, "Quantitative analysis of aromatic hydrocarbons by capillary gas chromatography", *ANAL. CHEM.* **35** (1963) 2022–2027.
- [16] H. Miyake, M. Mitooka, T. Matsumoto, "Studies of gas chromatography with mixed stationary liquids. Part IV. The analysis of aromatic hydrocarbons from C₆ to C₁₀ with a capillary column", *BUL. CHEM. SOC. JAPAN* **38** (1965) 1062–1071.
- [17] D.E. Willis, R.M. Engelbrecht, "Gas-chromatographic analysis of C₁ to C₁₀ hydrocarbons by open tubular columns with on-column injection", *J. GAS CHROMATOGR.* **5** (10) (1967) 536–538 (CA 68:41814r).
- [18] E. Matisova, "Identification of alkylbenzenes by capillary gas chromatography and combined gas chromatography mass spectrometry", *J. CHROMATOGR.* **438** (1988) 131–144.
- [19] E. Matisova, M. Rukriglova, J. Krupcik, E. Kovacicova, S. Holotik, "Identification of alkylbenzenes up to C₁₂ by capillary gas chromatography and combined gas chromatography-mass spectrometry", *J. CHROMATOGR.* **455** (1988) 301–309.
- [20] E. Matisova, E. Kovacicova, P.T. Ha, E. Kolek, W. Engewald, "Identification of alkylbenzenes up to C₁₂ by capillary gas chromatography and gas chromatography mass spectrometry. Part II. Retention indices on OV-101 columns and retention-molecular structure correlations", *J. CHROMATOGR.* **475** (1989) 113–123.
- [21] E. Matisova, P. Kuran, "Reproducibility of retention indices measurement of alkylbenzenes on cross-linked methyl silicone phase fused silica capillaries", *CHROMATOGRAPHY* **30** (1990) 328–332.
- [22] N. Dimov, E. Matisova, "Improved method for prediction of gas chromatographic retention indices of C₉–C₁₂ alkylbenzenes", *J. CHROMATOGR.* **549** (1991) 325–333.
- [23] E. Matisova, E. Juranyiova, P. Kuran, E. Brandsteterova, A. Kocan, S. Holotik, "Analysis of multi-component mixtures by high-resolution capillary gas chromatography and combined gas chromatography–mass spectrometry. Part I. Aromatics in a hydrocarbon matrix", *J. CHROMATOGR.* **552** (1991) 301–312.
- [24] E. Matisova, L. Kubus, E. Juranyiova, "Determination of traces of aromatic compounds in multi-component mixtures by HRCGC", *J. HIGH RESOLUT. CHROMATOGR.* **14** (1991) 713–717.
- [25] E. Matisova, "High resolution capillary gas chromatography of aromatic compounds in multi-component hydrocarbon mixtures", *J. HIGH RESOLUT. CHROMATOGR.* **15** (1992) 213–218.
- [26] R. Kaiser, "New developments in gas chromatography from the 1961 literature", *Z. ANAL. CHEM.* **189** (1962) 1–14 (CA 57:13163e).

Hivatkozások a 4. szakaszban

- [1] T. Tóth, F. Garay, Cs. Lovász, "A simple coupling device for flow tuning of serially coupled flexible fused silica capillary columns", J. HIGH RESOLUT. CHROMATOGR. **13** (1990) 840–842.

Hivatkozások a 6. szakaszban

- [1] "Nomenclature for chromatography (IUPAC recommendations 1993)", prepared for publication by L.S. Etre, PURE APPL. CHEM. **65** (1993) 819–872.
- [2] "Manual of symbols and terminology for physicochemical quantities and units (Adopted by the IUPAC Council on 7 July 1969)", prepared for publication by M.L. McGlashan, revision prepared for publication by M.A. Paul, second revision prepared for publication by D.H. Whiffen, PURE APPL. CHEM. **51** (1979) 1–41.
- [3] W. Kimpenshaus, F. Richter, L. Rohrschneider, CHROMATOGRAPHIA **15** (1982) 577.
- [4] L.S. Ettre, "Viscosity of gases used as the mobile phases in gas chromatography", CHROMATOGRAPHIA **18** (1984) 243–248.
- [5] S.J. Hawkes, "Viscosities of carrier gases at gas chromatograph temperatures and pressures", CHROMATOGRAPHIA **37** (1993) 399–401.
- [6] W. Licht, D.G. Stechert, "The variation of the viscosity of gases and vapors with temperature", J. PHYS. CHEM. **48** (1944) 23–47.
- [7] Valkó P, Vajda S., „Műszaki tudományos feladatok megoldása személyi számítógéppel”, Műszaki Könyvkiadó, Budapest (1987) 166. oldal.
- [8] Y.S. Touloukian, S.C. Saxena, P. Hestermans, Eds., "Thermophysical properties of matter", Vol. 11, IFI/Plenum, New York, Washington (1975).